

## Synaptophysin1 (p38-1)

Cat.No. 101 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> yes <b>ICC:</b> 1 : 1000 <b>IHC:</b> 1 : 500 <b>IHC-P (FFPE):</b> 1 : 200 <b>iDISCO:</b> external data (see remarks) <b>Clarity:</b> external data (see remarks)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of human Synaptophysin (UniProt Id: P08247)
Reactivity	Reacts with: human (P08247), rat (P07825), mouse (Q62277), hamster, cow, chicken, frog. Other species not tested yet.
Matching control	101-0P
Remarks	<b>iDISCO:</b> This antibody has been successfully applied and published for this method by customers (see application-specific references). <b>Clarity:</b> This antibody has been successfully applied and published for this method by customers (see application-specific references).

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

**Synaptophysin1**, also referred to as **p38-1**, is a membrane glycoprotein of synaptic vesicles that is ubiquitously expressed in all neurons and in many endocrine cells. It is currently the most widely used marker for nerve terminals and probably the best marker for the pathologist in differentiating neuroendocrine tumors.

Synaptophysin1 has four transmembrane domains with both N- and C-terminus facing the cytoplasm. It binds to synaptobrevin1 and synaptobrevin2 in detergent extracts but its function has not been elucidated completely. It forms a complex with dynamin at high Ca<sup>2+</sup> concentration suggesting an involvement in synaptic vesicle endocytosis. As typical for synaptic vesicle proteins, synaptophysin1 represents a small protein family with two additional members, synaptoporin (synaptophysin2) and panthophysin. Like synaptophysin1, synaptoporin is widely expressed in neurons and colocalizes with synaptophysin1 on synaptic vesicles whereas panthophysin is expressed in all tissues.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

### Selected References for 101 004

- Plekhh5-regulated autophagy of synaptic vesicles reveals a pathogenic mechanism in motoneuron disease. Lüningschrör P, Binotti B, Dombert B, Heimann P, Perez-Lara A, Slotta C, Thau-Habermann N, R von Collenberg C, Karl F, Damme M, Horowitz A, et al. Nature communications (2017) 81: 678. . **IHC, ICC, WB**
- Regulatory Function of Sympathetic Innervation on the Endo/Lysosomal Trafficking of Acetylcholine Receptor. Straka T, Schröder C, Roos A, Kollipara L, Sickmann A, Williams MPI, Hafner M, Khan MM, Rudolf R. Frontiers in physiology (2021) 12: 626707. . **IHC, iDISCO; tested species: mouse**
- Synaptophysin density underestimates synapse counts in the human brain - An anatomical study on synaptoporin protein expression. Woelfle S, Schön M, Pedro MT, Wagner J, Shiban E, Hong B, Tsoy Y, Boeckers TM. Neurobiology of disease (2026) 227: 107503. . **WB, IHC; tested species: human**
- Loss of synaptic Munc13-1 underlies neurotransmission abnormalities in spinal muscular atrophy. Moradi M, Deng C, Sendtner M. Cellular and molecular life sciences : CMLS (2025) 821: 325. . **WB, IHC; tested species: mouse**
- Manipulation of DHPS activity affects dendritic morphology and expression of synaptic proteins in primary rat cortical neurons. Cavalli P, Raffauf A, Passarella S, Helmuth M, Dieterich DC, Landgraf P. Frontiers in cellular neuroscience (2024) 18: 1465011. . **WB, ICC; tested species: rat**
- The lncRNA Malat1 is trafficked to the cytoplasm as a localized mRNA encoding a small peptide in neurons. Xiao W, Halabi R, Lin CH, Nazim M, Yeom KH, Black DL. Genes & development (2024) : . . **WB, ICC; tested species: mouse**
- Plastin 3 rescues cell surface translocation and activation of TrkB in spinal muscular atrophy. Hennlein L, Ghanawi H, Gerstner F, Palominos García E, Yildirim E, Saal-Bauernschubert L, Moradi M, Deng C, Klein T, Appenzeller S, Sauer M, et al. The Journal of cell biology (2023) 2223: . . **ICC, IHC; tested species: mouse**
- Synaptic vesicle proteins are selectively delivered to axons in mammalian neurons. Watson ET, Pauers MM, Seibert MJ, Vevea JD, Chapman ER. eLife (2023) 12: . . **ICC, WB**
- Astrocytic TDP-43 dysregulation impairs memory by modulating antiviral pathways and interferon-inducible chemokines. Licht-Murava A, Meadows SM, Palaguachi F, Song SC, Jackvony S, Bram Y, Zhou C, Schwartz RE, Froemke RC, Orr AL, Orr AG, et al. Science advances (2023) 916: eade1282. . **ICC, IHC; tested species: mouse**

Access the online factsheet including applicable protocols at <https://sysy.com/product/101004> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.