

## Synaptophysin1

Cat.No. 101 011C3; Monoclonal mouse antibody, 50 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg purified IgG, lyophilized, fluorescence-labeled with Cyanine 3. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Either add 1:1 (v/v) glycerol, then aliquot and store at -20°C until use, or store aliquots at -80°C without additives. Reconstitute immediately upon receipt! Avoid bright light when working with the antibody to minimize photo bleaching of the fluorescent dye. For detailed information, see back of the data sheet.
Applications	<b>WB:</b> N/A <b>IP:</b> N/A <b>ICC:</b> 1 : 100 up to 1 : 1000 <b>IHC:</b> 1 : 500 up to 1 : 1000 <b>IHC-P:</b> not tested yet
Label	Sulfo-Cyanine 3
Clone	7.2
Subtype	IgG1 (λ light chain)
Immunogen	Full-length recombinant rat synaptophysin (UniProt Id: P07825)
Epitop	AA 219 to 307 from rat Synaptophysin1 (UniProt Id: P07825)
Reactivity	Reacts with: human (P08247), rat (P07825), mouse (Q62277), mammals. Weaker signal: zebrafish, other vertebrates. Other species not tested yet.
Specificity	K.O. validated
Remarks	Widely used as marker for nerve terminals and neuroendocrine tumors. For unknown reason, neuronal synaptophysin is better recognised than neuroendocrine synaptophysin. If this is a problem, the polyclonal rabbit antibody, cat. no. 101 002 or 101 203 are recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Synaptophysin1**, also referred to as **p38-1**, is a membrane glycoprotein of synaptic vesicles that is ubiquitously expressed in all neurons and in many endocrine cells. It is currently the most widely used marker for nerve terminals and probably the best marker for the pathologist in differentiating neuroendocrine tumors.

Synaptophysin1 has four transmembrane domains with both N- and C-terminus facing the cytoplasm. It binds to synaptobrevin1 and synaptobrevin2 in detergent extracts but its function has not been elucidated completely. It forms a complex with dynamin at high Ca<sup>2+</sup> concentration suggesting an involvement in synaptic vesicle endocytosis. As typical for synaptic vesicle proteins, synaptophysin1 represents a small protein family with two additional members, synaptoporin (synaptophysin2) and panthophysin. Like synaptophysin1, synaptoporin is widely expressed in neurons and colocalizes with synaptophysin1 on synaptic vesicles whereas panthophysin is expressed in all tissues.

## Selected References for 101 011C3

Tau hyperphosphorylation in synaptosomes and neuroinflammation are associated with canine cognitive impairment. Smolek T, Madari A, Farbakova J, Kandrak O, Jadhav S, Cente M, Brezovakova V, Novak M, Zilka N. The Journal of comparative neurology (2016) 5244: 874-95. . **WB**

Impaired excitatory drive to spinal GABAergic neurons of neuropathic mice. Leitner J, Westerholz S, Heinke B, Forsthuber L, Wunderbaldinger G, Jäger T, Gruber-Schoffnegger D, Braun K, Sandkühler J. PLoS one (2013) 88: e73370. . **IHC; tested species: mouse**

Synapse formation and function is modulated by the amyloid precursor protein. Priller C, Bauer T, Mitteregger G, Krebs B, Kretschmar HA, Herms J. The Journal of neuroscience : the official journal of the Society for Neuroscience (2006) 2627: 7212-21. . **ICC**

Alterations in neuromuscular junction morphology with ageing and endurance training modulate neuromuscular transmission and myofibre composition. Yamaguchi T, Kouzaki K, Sasaki K, Nakazato K. The Journal of physiology (2024) : . . **IHC; tested species: mouse**

Conditioning- and reward-related dendritic and presynaptic plasticity of nucleus accumbens neurons in male and female sign-tracker rats. Colom M, Kraev I, Stramek AK, Loza IB, Rostron CL, Heath CJ, Dommett EJ, Singer BF. The European journal of neuroscience (2024) 607: 5694-5717. . **IHC; tested species: rat**

The activation of mGluR4 rescues parallel fiber synaptic transmission and LTP, motor learning and social behavior in a mouse model of Fragile X Syndrome. Martín R, Suárez-Pinilla AS, García-Font N, Laguna-Luque ML, López-Ramos JC, Oset-Gasque MJ, Gruart A, Delgado-García JM, Torres M, Sánchez-Prieto J. Molecular autism (2023) 141: 14. . **ICC; tested species: mouse**

Selective endocytosis of Ca<sup>2+</sup>-permeable AMPARs by the Alzheimer's disease risk factor CALM bidirectionally controls synaptic plasticity. Azarnia Tehran D, Kochlamazashvili G, Pampaloni NP, Sposini S, Shergill JK, Lehmann M, Pashkova N, Schmidt C, Löwe D, Napieczynska H, Heuser A, et al. Science advances (2022) 821: eabl5032. . **WB; tested species: mouse**

Transient oxytocin signaling primes the development and function of excitatory hippocampal neurons. Ripamonti S, Ambroziewicz MC, Guzzi F, Gravati M, Biella G, Bormuth I, Hammer M, Tuffy LP, Sigler A, Kawabe H, Nishimori K, et al. eLife (2017) 6: . . **ICC; tested species: mouse**

## Selected General References

Essential roles in synaptic plasticity for synaptogyrin I and synaptophysin I. Janz R et al. Neuron (1999) PubMed:10595519

Access the online factsheet including applicable protocols at <https://sysy.com/product/101011C3> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.