

Synaptoporin (p38-2)

Cat.No. 102 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 500 IHC: 1 : 1000 up to 1 : 2000 (see remarks) IHC-P (FFPE): 1 : 1000
Clone	44E3D12
Subtype	IgG2b (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat Synaptoporin (UniProt Id: P22831)
Reactivity	Reacts with: mouse (Q8BGN8), rat (P22831). Other species not tested yet.
Matching control	102-1P
Remarks	IHC: In addition to the expected strong mossy fiber staining, this antibody shows some weak unspecific staining of cell bodies in the cortex, striatum and cerebellum.

Background

Synaptoporin, also known as **synaptophysin2** and **p38-2**, is highly homologous to synaptophysin1 but encoded by a different gene. Like synaptophysin1, synaptoporin contains four transmembrane regions and a short cytoplasmic tail. Unlike synaptophysin1, it is not glycosylated. The distributions of synaptophysin1 and synaptoporin are different. Synaptophysin1 is more uniformly expressed whereas synaptoporin is particularly enriched in mossy fiber synapses in the hippocampus. It is thus an excellent marker for subsets of synapses.

Selected References for 102 011

Mossy Fiber Sprouting in Temporal Lobe Epilepsy: The Impact of Netrin-1, DCC, and Gene Expression Changes. Onay M, Harter PN, Weber K, Piiper A, Czabanka M, Plate KH, Freiman TM, Gessler F, Puhahn-Schmeiser B Biomedicines (2024) 1212: . . . **IHC-P; tested species: human**

Selected General References

The synaptic vesicle cycle: a cascade of protein-protein interactions. Südhof TC et al. Nature (1995) PubMed:7791897

Synaptic vesicles and exocytosis. Jahn R et al. Annu. Rev. Neurosci. (1994) PubMed:8210174

Relative properties and localizations of synaptic vesicle protein isoforms: the case of the synaptophysins. Fykse EM et al. J. Neurosci. (1993) PubMed:8229211

Synaptoporin, a novel putative channel protein of synaptic vesicles. Knaus P et al. Neuron (1990) PubMed:2206533

Access the online factsheet including applicable protocols at <https://susy.com/product/102011> or scan the QR-code.



TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.