

Synaptotagmin7

Cat.No. 105-71P; control protein, 100 µg protein (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg protein, lyophilized. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in TBS. Then aliquot and store at -20°C to -80°C until use. Control proteins should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Immunogen	Recombinant protein corresponding to AA 46 to 133 from rat Synaptotagmin7 (UniProt Id: Q62747)
Recommended dilution	Optimal concentrations should be determined by the end-user.
Matching antibodies	105 173
Remarks	This control protein consists of the Strep-Tag® fusion protein (aa 46 - 133 of rat synaptotagmin 7) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Synaptotagmin7 is a proposed regulator of Ca²⁺ dependent exocytosis like neurotransmitter release. It occurs in several splicing variants which are expressed in a developmentally regulated pattern in brain. The distinct roles for the alternative splicing isoforms have not yet been determined. Synaptotagmin7 shows Ca²⁺ dependent oligomerization via its own C2 domains leading to the formation of large linear structures which reside at the fusion site of vesicles and plasma membrane. These oligomers may be involved in the modulation of Ca²⁺ dependent exocytosis by opening or dilating fusion pores.

Selected General References

Alternative splicing isoforms of synaptotagmin VII in the mouse, rat and human.
Fukuda M, Ogata Y, Saegusa C, Kanno E, Mikoshiba K
The Biochemical journal (2002) 365Pt 1: 173-80. .

The calcium-binding loops of the tandem C2 domains of synaptotagmin VII cooperatively mediate calcium-dependent oligomerization.

Fukuda M, Katayama E, Mikoshiba K
The Journal of biological chemistry (2002) 27732: 29315-20. .

Synaptotagmin VII as a plasma membrane Ca(2+) sensor in exocytosis.
Sugita S, Han W, Butz S, Liu X, Fernández-Chacón R, Lao Y, Südhof TC
Neuron (2001) 302: 459-73. .

Mechanism of the calcium-dependent multimerization of synaptotagmin VII mediated by its first and second C2 domains.
Fukuda M, Mikoshiba K
The Journal of biological chemistry (2001) 27629: 27670-6. .

The Exocytosis-regulatory protein synaptotagmin VII mediates cell invasion by Trypanosoma cruzi.
Caler EV, Chakrabarti S, Fowler KT, Rao S, Andrews NW
The Journal of experimental medicine (2001) 1939: 1097-104. .

Distinct self-oligomerization activities of synaptotagmin family. Unique calcium-dependent oligomerization properties of synaptotagmin VII.

Fukuda M, Mikoshiba K
The Journal of biological chemistry (2000) 27536: 28180-5. .

Access the online factsheet including applicable protocols at <https://susy.com/product/105-71P> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.