

## Rab3a

Cat.No. 107 102; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For <b>reconstitution</b> add 200 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> yes <b>ICC:</b> 1 : 1000 <b>IHC:</b> yes <b>IHC-P:</b> not tested yet <b>ELISA:</b> yes (see remarks)
Immunogen	Synthetic peptide corresponding to AA 2 to 14 from rat Rab3a (UniProt Id: P63012)
Reactivity	Reacts with: human (P20336), rat (P63012), mouse (P63011), hamster, cow, zebrafish. Other species not tested yet.
Matching control	107-1P
Remarks	<b>ELISA:</b> Suitable as detector antibody for sandwich-ELISA with cat. no. <a href="#">107 111</a> as capture antibody. The ELISA-protocol for membrane proteins is recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Rab 3** is a member of the Rab protein family that belongs to the ras-related superfamily of small monomeric GTPases. Four related isoforms of Rab 3 are known (**Rab 3a**, **3b**, **3c**, and **3d**). Rab 3a and 3c are predominantly expressed in neurons and neuroendocrine cells where they are localized to synaptic vesicles. Unlike the integral membrane proteins of synaptic vesicles, Rab 3a/c is absent from the Golgi complex and thus does not result in immunostaining of the axo-dendritic region as sometimes seen with e.g. synaptophysin, synaptobrevin/VAMP, or synaptogyrin. Rab 3b and 3d are expressed in non-neuronal tissues such as adipocytes and the exocrine pancreas (3d). It has been shown that overexpression of Rab 3 inhibits Ca<sup>2+</sup> regulated exocytosis and converts it into an constitutive Ca<sup>2+</sup> independent exocytosis mechanism.

## Selected References for 107 102

Localization versus function of Rab3 proteins. Evidence for a common regulatory role in controlling fusion. Schlüter OM, Khvotchev M, Jahn R, Südhof TC  
The Journal of biological chemistry (2002) 27743: 40919-29. . **WB, IHC**

SV31 is a Zn<sup>2+</sup>-binding synaptic vesicle protein. Barth J, Zimmermann H, Volkandt W  
Journal of neurochemistry (2011) 1184: 558-70. . **WB, ICC**

Regulation of density of functional presynaptic terminals by local energy supply. Zhou H, Liu G  
Molecular brain (2015) 8: 42. . **WB, ICC**

Distribution of synaptic vesicle proteins in the mammalian retina identifies obligatory and facultative components of ribbon synapses. Von Kriegstein K, Schmitz F, Link E, Südhof TC  
The European journal of neuroscience (1999) 114: 1335-48. . **WB, IHC**

Synapsin-dependent reserve pool of synaptic vesicles supports replenishment of the readily releasable pool under intense synaptic transmission. Vasileva M, Horstmann H, Geumann C, Gitler D, Kuner T  
The European journal of neuroscience (2012) 368: 3005-20. . **ELISA**

Cochlear ablation in neonatal rats disrupts inhibitory transmission in the medial nucleus of the trapezoid body. Hruskova B, Trojanova J, Kralikova M, Melichar A, Suchankova S, Bartosova J, Burianova JS, Popelar J, Syka J, Turecek R  
Neuroscience letters (2019) : . . **IHC; tested species: rat**

The GTPase Rab26 links synaptic vesicles to the autophagy pathway. Binotti B, Pavlos NJ, Riedel D, Wenzel D, Vorbrüggen G, Schalk AM, Kühnel K, Boyken J, Erck C, Martens H, Chua JJ, et al.  
eLife (2015) 4: e05597. . **WB**

Proteomic screening of glutamatergic mouse brain synaptosomes isolated by fluorescence activated sorting. Biesemann C, Grønberg M, Luquet E, Wichert SP, Bernard V, Bungers SR, Cooper B, Varoquaux F, Li L, Byrne JA, Urlaub H, et al.  
The EMBO journal (2014) 332: 157-70. . **WB; tested species: mouse**

Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins. Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rammner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, et al.  
Science (New York, N.Y.) (2014) 3446187: 1023-8. . **IHC; tested species: mouse**

Glomerular podocytes possess the synaptic vesicle molecule Rab3A and its specific effector rabphilin-3a. Rastaldi MP, Armelloni S, Berra S, Li M, Pesaresi M, Poczewski H, Langer B, Kerjaschki D, Henger A, Blattner SM, Kretzler M, et al.  
The American journal of pathology (2003) 1633: 889-99. . **IHC**

Access the online factsheet including applicable protocols at <https://sysy.com/product/107102> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.