

## Rab3a

Cat.No. 107 111; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> yes <b>ICC:</b> 1 : 1000 up to 1 : 2000 <b>IHC:</b> 1 : 500 <b>IHC_P:</b> 1 : 2000 up to 1 : 3000 <b>DNA_PAINT:</b> yes (see remarks) <b>EM:</b> yes <b>ELISA:</b> yes (see remarks)
Clone	42.2
Subtype	IgG1 (κ light chain)
Immunogen	Full length rat recombinant Rab3a (UniProt Id: P63012)
Epitop	Epitop: AA 191 to 220 from rat Rab3a (UniProt Id: P63012)
Reactivity	Reacts with: human (P20336), rat (P63012), mouse (P63011), mammals. Other species not tested yet.
Specificity	K.O. PubMed: <a href="https://pubmed.ncbi.nlm.nih.gov/12244319/">12244319</a>
Remarks	<b>DNA_PAINT:</b> This antibody has been successfully used for DNA-PAINT application (see Unterauer et al., 2024; PMID: 38552614). <b>ELISA:</b> Suitable as capture antibody for sandwich-ELISA with cat. no. 107 102 as detector antibody. The ELISA-protocol for membrane proteins is recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Rab 3** is a member of the Rab protein family that belongs to the ras-related superfamily of small monomeric GTPases. Four related isoforms of Rab 3 are known (**Rab 3a**, **3b**, **3c**, and **3d**). Rab 3a and 3c are predominantly expressed in neurons and neuroendocrine cells where they are localized to synaptic vesicles. Unlike the integral membrane proteins of synaptic vesicles, Rab 3a/c is absent from the Golgi complex and thus does not result in immunostaining of the axo-dendritic region as sometimes seen with e.g. synaptophysin, synaptobrevin/VAMP, or synaptogyrin. Rab 3b and 3d are expressed in non-neuronal tissues such as adipocytes and the exocrine pancreas (3d). It has been shown that overexpression of Rab 3 inhibits Ca<sup>2+</sup> regulated exocytosis and converts it into an constitutive Ca<sup>2+</sup> independent exocytosis mechanism.

## Selected References for 107 111

- Association of Rab3A with synaptic vesicles at late stages of the secretory pathway.  
Matteoli M, Takei K, Cameron R, Hurlbut P, Johnston PA, Südhof TC, Jahn R, De Camilli P  
The Journal of cell biology (1991) 1153: 625-33. . **ICC, IHC, WB, EM; tested species: rat, frog**
- Quantitative analysis of synaptic vesicle Rabs uncovers distinct yet overlapping roles for Rab3a and Rab27b in Ca<sup>2+</sup>-triggered exocytosis.  
Pavlos NJ, Grønberg M, Riedel D, Chua JJ, Boyken J, Kloeppe TH, Urlaub H, Rizzoli SO, Jahn R  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2010) 3040: 13441-53. . **WB, IP, ICC**
- Actions of Rab27B-GTPase on mammalian central excitatory synaptic transmission.  
Arias-Hervert ER, Xu N, Njus M, Murphy GG, Hou Y, Williams JA, Lentz SI, Ernst SA, Stuenkel EL  
Physiological reports (2020) 89: e14428. . **WB, IHC; tested species: mouse**
- JIP3 localises to exocytic vesicles and focal adhesions in the growth cones of differentiated PC12 cells.  
Caswell PT, Dickens M  
Molecular and cellular biochemistry (2017) : . . **WB, ICC; tested species: rat**
- BDNF enhances spontaneous and activity-dependent neurotransmitter release at excitatory terminals but not at inhibitory terminals in hippocampal neurons.  
Shinoda Y, Ahmed S, Ramachandran B, Bharat V, Brockelt D, Altas B, Dean C  
Frontiers in synaptic neuroscience (2014) 6: 27. . **WB, ICC; tested species: rat**
- Myosin5a tail associates directly with Rab3A-containing compartments in neurons.  
Wöllert T, Patel A, Lee YL, Provance DW, Vought VE, Cosgrove MS, Mercer JA, Langford GM  
The Journal of biological chemistry (2011) 28616: 14352-61. . **WB, ICC; tested species: mouse**
- Biochemical, molecular and behavioral phenotypes of Rab3A mutations in the mouse.  
Yang S, Farias M, Kapfhamer D, Tobias J, Grant G, Abel T, Bučan M  
Genes, brain, and behavior (2007) 61: 77-96. . **WB, IP**
- Rabphilin knock-out mice reveal that rabphilin is not required for rab3 function in regulating neurotransmitter release.  
Schlüter OM, Schnell E, Verhage M, Tzonopoulos T, Nicoll RA, Janz R, Malenka RC, Geppert M, Südhof TC  
The Journal of neuroscience : the official journal of the Society for Neuroscience (1999) 1914: 5834-46. . **WB, IHC**
- Synaptic targeting of rabphilin-3A, a synaptic vesicle Ca<sup>2+</sup>/phospholipid-binding protein, depends on rab3A/3C.  
Li C, Takei K, Geppert M, Daniell L, Stenius K, Chapman ER, Jahn R, De Camilli P, Südhof TC  
Neuron (1994) 134: 885-98. . **WB, IHC**
- Spatial proteomics in neurons at single-protein resolution.  
Unterauer EM, Shetab Boushehri S, Jevdokimenko K, Masullo LA, Ganji M, Sograte-Idrissi S, Kowalewski R, Strauss S, Reinhardt SCM, Perovic A, Marr C, et al.  
Cell (2024) 1877: 1785-1800.e16. . **DNA\_PAINT; tested species: rat**

Access the online factsheet including applicable protocols at <https://sysy.com/product/107111> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.