

## Syntaxin3

Cat.No. 110 032; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/ Storage	200 µl antiserum, lyophilized. For <b>reconstitution</b> add 200 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) (see remarks) <b>IP:</b> yes <b>ICC:</b> 1 : 100 <b>IHC:</b> 1 : 2500 <b>IHC-P:</b> not tested yet
Immunogen	Recombinant protein corresponding to AA 1 to 260 from rat Syntaxin3 (UniProt ID: Q08849)
Reactivity	Reacts with: human (Q13277), rat (Q08849), mouse (Q64704), hamster, pig, zebrafish. Other species not tested yet.
Matching control	110-3P
Remarks	<b>WB:</b> The affinity purified antibody (cat. no. 110 033) gives stronger signals in Western blots.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Syntaxin 3**, a member of the SNARE family of proteins, is related to syntaxin 1. While often co-expressed with syntaxins 2 and 4 within the same cell type, its membrane localization is usually different, for instance in epithelial cells or in the exocrine pancreas. Like syntaxins 1, 2 and 4 it appears to be involved in the fusion of transport vesicles with the plasma membrane.

## Selected References for 110 032

- Sunday driver interacts with two distinct classes of axonal organelles.  
Abe N, Almenar-Queralt A, Lillo C, Shen Z, Lozach J, Briggs SP, Williams DS, Goldstein LS, Cavalli V  
The Journal of biological chemistry (2009) 28450: 34628-39. . **WB, IP, IHC**
- SNAREs Interact with Retinal Degeneration Slow and Rod Outer Segment Membrane Protein-1 during Conventional and Unconventional Outer Segment Targeting.  
Zulliger R, Conley SM, Mwoyosvi ML, Stuck MW, Azadi S, Naash MI  
PloS one (2015) 109: e0138508. . **WB, IP**
- Axonal BACE1 dynamics and targeting in hippocampal neurons: a role for Rab11 GTPase.  
Buggia-Prévoit V, Fernandez CG, Riordan S, Vetrivel KS, Roseman J, Waters J, Bindokas VP, Vassar R, Thinakaran G  
Molecular neurodegeneration (2014) 9: 1. . **ICC**
- The stability of the primed pool of synaptic vesicles and the clamping of spontaneous neurotransmitter release rely on the integrity of the C-terminal half of the SNARE domain of syntaxin-1A.  
Salazar Lázaro A, Trimbuch T, Vardar G, Rosenmund C  
eLife (2024) 12: . . **WB; tested species: mouse**
- Knockout of Syntaxin-4 in 3T3-L1 adipocytes reveals new insight into GLUT4 trafficking and adiponectin secretion.  
Black HL, Livingstone R, Mastick CC, Al Tobi M, Taylor H, Geiser A, Stirrat L, Kioumourtzoglou D, Petrie JR, Boyle JG, Bryant NJ, et al.  
Journal of cell science (2021) : . . **WB; tested species: mouse**
- Cardiac SNARE Expression in Health and Disease.  
Bowman PRT, Smith GL, Gould GW  
Frontiers in endocrinology (2019) 10: 881. . **WB; tested species: mouse**
- Identification of a Botulinum Neurotoxin-like Toxin in a Commensal Strain of Enterococcus faecium.  
Zhang S, Lebreton F, Mansfield MJ, Miyashita SI, Zhang J, Schwartzman JA, Tao L, Masuyer G, Martínez-Carranza M, Stenmark P, Gilmore MS, et al.  
Cell host & microbe (2018) 232: 169-176.e6. . **WB; tested species: mouse**
- Vesicle associated membrane protein 8 (VAMP8)-mediated zymogen granule exocytosis is dependent on endosomal trafficking via the constitutive-like secretory pathway.  
Messenger SW, Falkowski MA, Thomas DD, Jones EK, Hong W, Gaisano HY, Giasano HY, Boullis NM, Groblewski GE  
The Journal of biological chemistry (2014) 28940: 28040-53. . **WB**
- Tumor protein D52 controls trafficking of an apical endolysosomal secretory pathway in pancreatic acinar cells.  
Messenger SW, Thomas DD, Falkowski MA, Byrne JA, Gorelick FS, Groblewski GE  
American journal of physiology. Gastrointestinal and liver physiology (2013) 3056: G439-52. . **WB; tested species: rat**
- Myosin Ia is required for CFTR brush border membrane trafficking and ion transport in the mouse small intestine.  
Kravtsov DV, Caputo C, Collaco A, Hoekstra N, Egan ME, Mooseker MS, Ameen NA  
Traffic (Copenhagen, Denmark) (2012) 138: 1072-82. . **IHC; tested species: mouse**
- How pig sperm prepares to fertilize: stable acrosome docking to the plasma membrane.  
Tsai PS, Garcia-Gil N, van Haeften T, Gadella BM  
PloS one (2010) 56: e11204. . **WB; tested species: pig**

Access the online factsheet including applicable protocols at <https://sysy.com/product/110032> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.