

Syntaxin1A

Cat.No. 110 318; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP-staining) IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 (see remarks) IHC_P: 1 : 1000
Clone	Gp78.3
Subtype	IgG2 (κ light chain)
Immunogen	Recombinant protein corresponding to cytoplasmic-domain without membrane anchor of rat Syntaxin 1A. (UniProt Id: P32851)
Reactivity	Reacts with: human (Q16623), rat (P32851), mouse (O35526), mammals, chicken. Other species not tested yet.
Specificity	Specific für syntaxin 1A, no cross-reactivity to syntaxin 1B. K.O.
Matching control	110-1P
Remarks	This antibody is a chimeric antibody based on the monoclonal mouse antibody clone 78.3. The constant regions of the heavy and light chains have been replaced by guinea pig specific sequences. Therefore, the antibody can be used with standard anti-guinea pig secondary reagents. The antibody has been expressed in mammalian cells. IHC: Heat-mediated antigen retrieval (in citrate buffer pH 6) is required for immunohistochemical staining.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Syntaxin 1, also known as **p35**, is a small integral membrane protein that is abundantly expressed in neurons and neuroendocrine cells. It was initially discovered as HPC-1. Syntaxin 1 is an essential component of the exocytotic fusion machine and interacts with several other proteins important for synaptic function, including its partners in the fusion complex synaptobrevin, SNAP 25, α-SNAP, synaptotagmin 1, Munc 18/n-Sec1 and Ca²⁺-channels.

Syntaxin 1 is localized primarily to the neuronal plasmalemma and is concentrated in synapses where pools of the protein are also present on recycling organelles including synaptic vesicles. It is the main target of one of the Botulinum neurotoxins BoNT/C1 which, however, cannot cleave the protein when complexed with its partner proteins in the fusion complex.

Selected General References

Mechanisms of synaptic vesicle exocytosis.

Lin RC, Scheller RH

Annual review of cell and developmental biology (2000) 16: 19-49. .

Phosphorylated syntaxin 1 is localized to discrete domains along a subset of axons.

Foletti DL, Lin R, Finley MA, Scheller RH

The Journal of neuroscience : the official journal of the Society for Neuroscience (2000) 2012: 4535-44. .

Membrane fusion and exocytosis.

Jahn R, Südhof TC

Annual review of biochemistry (1999) 68: 863-911. .

The synaptic vesicle cycle: a cascade of protein-protein interactions.

Südhof TC

Nature (1995) 3756533: 645-53. .

Synaptic vesicles and exocytosis.

Jahn R, Südhof TC

Annual review of neuroscience (1994) 17: 219-46. .

Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones.

Bennett MK, Calakos N, Scheller RH

Science (New York, N.Y.) (1992) 2575067: 255-9. .

Cloning and sequence analysis of cDNA for a neuronal cell membrane antigen, HPC-1.

Inoue A, Obata K, Akagawa K

The Journal of biological chemistry (1992) 26715: 10613-9. .

Epimorphin: a mesenchymal protein essential for epithelial morphogenesis.

Hirai Y, Takebe K, Takashina M, Kobayashi S, Takeichi M

Cell (1992) 693: 471-81. .

A marker of early amacrine cell development in rat retina.

Barnstable CJ, Hofstein R, Akagawa K

Brain research (1985) 3522: 286-90. .

Access the online factsheet including applicable protocols at <https://susy.com/product/110318> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.