

## $\alpha/\beta$ SNAP

Cat.No. 112 111; Monoclonal mouse antibody, 100  $\mu$ g purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 $\mu$ g purified IgG, lyophilized. For <b>reconstitution</b> add 100 $\mu$ l H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 up to 1 : 10000 (AP staining) <b>IP:</b> yes (see remarks) <b>ICC:</b> 1 : 500 up to 1 : 1000 <b>IHC:</b> external data (see remarks) <b>IHC-P (FFPE):</b> not tested yet
Clone	77.2
Subtype	IgG1 ( $\kappa$ light chain)
Immunogen	Recombinant protein corresponding to AA 1 to 295 from rat $\alpha$ SNAP (UniProt Id: P54921)
Reactivity	Reacts with: human (P54920, P60880), rat (P54921, P60881), mouse (Q9DB05, P28663), zebrafish. Other species not tested yet.
Specificity	Specific for $\alpha$ - and $\beta$ SNAP, does not cross-react to $\gamma$ SNAP.
Remarks	<b>IP:</b> The antibody does not immunoprecipitate the 20 S SNARE-complex. <b>IHC:</b> This antibody has been successfully applied and published for this method by customers (see application-specific references). It has not been validated using our standard protocols.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

The proteins  $\alpha/\beta$ -SNAP are two related soluble and highly conserved proteins that bind to the fusion complex (SNARE complex), thus allowing the N-ethylmaleimide sensitive fusion protein NSF to bind to the complex.  $\gamma$ -SNAP binds directly to NSF and Gaf-1/Rip11, a protein of the Rab11 interacting family. In contrast to  $\alpha/\beta$ -SNAP it does not interact directly with SNARE proteins and is not required for ER-Golgi transport. SNAP-proteins are abundantly expressed in all tissues. They are partially soluble, partially membrane-bound.

### Selected References for 112 111

- Composition of isolated synaptic boutons reveals the amounts of vesicle trafficking proteins. Wilhelm BG, Mandad S, Truckenbrodt S, Kröhnert K, Schäfer C, Rammner B, Koo SJ, Claßen GA, Krauss M, Haucke V, Urlaub H, et al. *Science (New York, N.Y.)* (2014) 3446187: 1023-8. . **WB, ICC, IHC; tested species: mouse, rat**
- Intersectin-Mediated Clearance of SNARE Complexes Is Required for Fast Neurotransmission. Jäpel M, Gerth F, Sakaba T, Bacetic J, Yao L, Koo SJ, Maritzen T, Freund C, Haucke V *Cell reports* (2020) 302: 409-420.e6. . **WB; tested species: mouse**
- Pleiotropic effects of alpha-SNAP M105I mutation on oocyte biology: ultrastructural and cellular changes that adversely affect female fertility in mice. de Paola M, Miró MP, Ratto M, Bätiz LF, Michaut MA *Scientific reports* (2019) 91: 17374. . **ICC; tested species: mouse**
- Cortical Granule Exocytosis Is Mediated by Alpha-SNAP and N-Ethylmaleimide Sensitive Factor in Mouse Oocytes. de Paola M, Bello OD, Michaut MA *PLoS one* (2015) 108: e0135679. . **WB**
- Ubiquitin-Synaptobrevin Fusion Protein Causes Degeneration of Presynaptic Motor Terminals in Mice. Liu Y, Li H, Sugiura Y, Han W, Gallardo G, Khvotchev M, Zhang Y, Kavalali ET, Südhof TC, Lin W *The Journal of neuroscience : the official journal of the Society for Neuroscience* (2015) 3533: 11514-31. . **WB**
- An essential and NSF independent role for  $\alpha$ -SNAP in store-operated calcium entry. Miao Y, Miner C, Zhang L, Hanson PI, Dani A, Vig M *eLife* (2013) 2: e00802. . **WB; KD verified**
- Doc2b is a high-affinity Ca<sup>2+</sup> sensor for spontaneous neurotransmitter release. Groffen AJ, Martens S, Diez Arazola R, Cornelisse LN, Lozovaya N, de Jong AP, Goriounova NA, Habets RL, Takai Y, Borst JG, Brose N, et al. *Science (New York, N.Y.)* (2010) 3275973: 1614-8. . **WB; tested species: mouse**
- alpha-SNAP and NSF are required in a priming step during the human sperm acrosome reaction. Tomes CN, De Blas GA, Michaut MA, Farré EV, Cherhiti O, Visconti PE, Mayorga LS *Molecular human reproduction* (2005) 111: 43-51. . **ICC; tested species: human**
- SNARE proteins are highly enriched in lipid rafts in PC12 cells: implications for the spatial control of exocytosis. Chamberlain LH, Burgoyne RD, Gould GW *Proceedings of the National Academy of Sciences of the United States of America* (2001) 9810: 5619-24. . **WB; tested species: rat**
- Comparison of cysteine string protein (Csp) and mutant alpha-SNAP overexpression reveals a role for csp in late steps of membrane fusion in dense-core granule exocytosis in adrenal chromaffin cells. Graham ME, Burgoyne RD *The Journal of neuroscience : the official journal of the Society for Neuroscience* (2000) 204: 1281-9. . **ICC**
- The N-ethylmaleimide-sensitive fusion protein and alpha-SNAP induce a conformational change in syntaxin. Hanson PI, Otto H, Barton N, Jahn R *The Journal of biological chemistry* (1995) 27028: 16955-61. . **WB**

Access the online factsheet including applicable protocols at <https://sysy.com/product/112111> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.