

## GluN1 extracellular

Cat.No. 114 018; Recombinant rabbit antibody, 50 µg recombinant IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> not recommended <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 (see remarks) <b>IHC:</b> 1 : 500 (see remarks) <b>IHC-P:</b> not recommended <b>IHC-G:</b> 1 : 500 (see remarks)
Clone	Rb003-102
Subtype	IgG1 (κ light chain)
Immunogen	Full length native human GluN1 (UniProt Id: Q05586)
Reactivity	Reacts with: mouse (P35438), rat (P35439). Other species not tested yet.
Remarks	This antibody is a chimeric antibody based on the monoclonal human antibody 003-102 (Kreye et al. 2016). The constant regions of the heavy and light chains have been replaced with rabbit specific sequences. The antibody can therefore be used with standard anti-rabbit secondary reagents. The antibody has been expressed in mammalian cells. <b>ICC:</b> This antibody is also suitable for the surface staining of living cells. After washing cells with bound antibodies, they can be fixed and visualized with secondary reagents. <b>IHC:</b> Antibody works on PFA/formaldehyde tissue. For optimal signal to noise ratio, glyoxal fixation (IHC-G) is recommended. <b>IHC-G:</b> 9% glyoxal, 8% acetic acid, in ddH <sub>2</sub> O, pH 4.2-4.4, according to Konno et al. 2023 is recommended.

## Background

**GluNs (NMDA-receptors)** represent a class of glutamate receptors that are of central importance in synaptic plasticity. Multiple NMDA receptor subtypes exist: **GluN1** and GluN2 A-D. GluN1 is the most important as it is required for activity. NMDA-receptors allow Ca<sup>2+</sup> influx and are thought to trigger Ca<sup>2+</sup> dependent postsynaptic processes involved in long term potentiation and depression.

## Selected General References

AMPA and NMDA receptors: similarities and differences in their synaptic distribution.  
Nusser Z et al. Curr. Opin. Neurobiol. (2000) PubMed:10851167

A nomenclature for ligand-gated ion channels.  
Collingridge GL et al. Neuropharmacology (2009) PubMed:18655795

Differential assembly of coexpressed glutamate receptor subunits in neurons of rat cerebral cortex.  
Brose N et al. J. Biol. Chem. (1994) PubMed:8207001

Molecular cloning and characterization of the rat NMDA receptor.  
Moriyoshi K et al. Nature (1991) PubMed:1834949

Access the online factsheet including applicable protocols at <https://sysy.com/product/114018> or scan the QR-code.



**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.