

## Munc13-1

Cat.No. 126 111; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 500 up to 1 : 5000 (AP staining) <b>IP:</b> yes (see remarks) <b>ICC:</b> not tested yet <b>IHC:</b> not tested yet <b>IHC-P:</b> not tested yet <b>ELISA:</b>
Clone	266B1
Subtype	IgG2b (κ light chain)
Immunogen	Recombinant protein corresponding to AA 3 to 317 from rat Munc13-1 (UniProt Id: Q62768)
Reactivity	Reacts with: rat (Q62768), mouse (Q4KUS2), zebrafish. Other species not tested yet.
Specificity	K.O. validated
Remarks	<b>IP:</b> For most effective IP, use the denaturing IP-protocol. Consider that protein-protein interactions may be affected.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Munc13s** are homologues of the *C. elegans* unc-13 gene product. Three brain-specific isoforms, Munc 13-1, -13-2, and -13-3, are expressed in rat, where they localize to presynaptic terminals. All three isoforms share multiple regulatory domains that may mediate phorbol ester and diacylglycerol binding.

**Munc13-1** shows the broadest expression pattern and is found in cortex, cerebellum, olfactory bulb, and hippocampus. Munc13-2 is mainly expressed in cortex and hippocampus, whereas **Munc13-3** exhibits highest expression levels in cerebellum and pons. Munc13-1 interacts directly with a putative coiled-coil domain in the N-terminal part of syntaxin and is involved in synaptic vesicle priming. For Munc13-2 an additional ubiquitously expressed N-terminal splice variant (ubMunc13-2) has been described.

Munc13-3 has been shown to be involved in the regulation of cerebellar synaptic transmission and motor learning.

## Selected References for 126 111

Formation of Golgi-derived active zone precursor vesicles.

Maas C, Torres VI, Altkrock WD, Leal-Ortiz S, Wagh D, Terry-Lorenzo RT, Fejtova A, Gundelfinger ED, Ziv NE, Garner CC  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2012) 3232: 11095-108. . **WB, ICC**

Different Munc13 isoforms function as priming factors in lytic granule release from murine cytotoxic T lymphocytes.

Dudenhöffer-Pfeifer M, Schirra C, Pattu V, Halimani M, Maier-Peuschel M, Marshall MR, Matti U, Becherer U, Dirks J, Jung M, Lipp P, et al.

Traffic (Copenhagen, Denmark) (2013) 147: 798-809. . **IP; tested species: mouse**

A presynaptic phosphosignaling hub for lasting homeostatic plasticity.

Müller JA, Betzin J, Santos-Tejedor J, Mayer A, Oprişoreanu AM, Engholm-Keller K, Paulußen I, Gulakova P, McGovern TD, Gschossman LJ, Schönhense E, et al.

Cell reports (2022) 393: 110696. . **WB; tested species: mouse**

CYP46A1 Activation by Efavirenz Leads to Behavioral Improvement without Significant Changes in Amyloid Plaque Load in the Brain of 5XFAD Mice.

Petrov AM, Lam M, Mast N, Moon J, Li Y, Maxfield E, Pikuleva IA

Neurotherapeutics : the journal of the American Society for Experimental NeuroTherapeutics (2019) : . . **WB; tested species: mouse**

RIM C2B Domains Target Presynaptic Active Zone Functions to PIP2-Containing Membranes.

de Jong APH, Roggero CM, Ho MR, Wong MY, Brautigam CA, Rizo J, Kaeser PS

Neuron (2018) 982: 335-349.e7. . **ICC; tested species: mouse**

Vti1a/b regulate synaptic vesicle and dense core vesicle secretion via protein sorting at the Golgi.

Emperador-Melero J, Huson V, van Weering J, Bollmann C, Fischer von Mollard G, Toonen RF, Verhage M

Nature communications (2018) 91: 3421. . **ICC; tested species: mouse**

RIM-binding protein 2 regulates release probability by fine-tuning calcium channel localization at murine hippocampal synapses.

Grauel MK, Maglione M, Reddy-Alla S, Willmes CG, Brockmann MM, Trimbuch T, Rosenmund T, Pangalos M, Vardar G, Stumpf A, Walter AM, et al.

Proceedings of the National Academy of Sciences of the United States of America (2016) 11341: 11615-11620. . **WB; tested species: mouse**

Synaptotagmin-12 phosphorylation by cAMP-dependent protein kinase is essential for hippocampal mossy fiber LTP.

Kaeser-Woo YJ, Younts TJ, Yang X, Zhou P, Wu D, Castillo PE, Südhof TC

The Journal of neuroscience : the official journal of the Society for Neuroscience (2013) 3323: 9769-80. . **WB**

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/126111> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.