SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

VGAT (SLC32A1) cytoplasmic domain

Cat.No. 131 011C3; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized, fluorescence-labeled with Cyanine 3. Albumin was added for stabilization. For **reconstitution** add 100 µl H_2O to get a 1mg/ml solution in TBS. Either add 1:1 (v/v) glycerol, then aliquot and store at -20°C until use, or store aliquots at -80°C without additives. Reconstitute immediately upon receipt! Avoid bright light when working with the antibody to minimize photo bleeching of the fluorescent dye. For detailed information, see back of the data sheet.
Applications	**WB**: N/A **IP**: N/A **ICC**: 1 : 1000 up to 1 : 2000 **IHC**: external data (see remarks) **IHC-P (FFPE)**: not tested yet
Label	Sulfo-Cyanine 3
Clone	117G4
Subtype	IgG3 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the amino terminus of rat VGAT (UniProt Id: O35458)
Reactivity	Reacts with: human (Q9H598), rat (O35458), mouse (O35633), Guinea pig, monkey. Other species not tested yet. Predicted to cross-react with cat due to high sequence homology.
Specificity	K.O. validated
Matching control	131-0P
Remarks	**IHC**: This antibody has been successfully applied and published for this method by customers (see application-specific references). It has not been validated using our standard protocols.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The **v**esicular **GA**BA **t**ransporter **VGAT** is responsible for uptake and storage of GABA and glycine by synaptic vesicles in the central nervous system. For this reason it is frequently referred to as the **v**esicular **i**nhibitory **a**mino**a**cid **t**ransporter **VIAAT**. It is different from the plasma membrane transporters in that it is driven by a proton electrochemical gradient across the vesicle membrane. So far, only one isoform is known. VGAT is currently the best marker for inhibitory nerve terminals.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

Selected References for 131 011C3

Cellular distribution of the NMDA-receptor activated synapto-nuclear messenger Jacob in the rat brain.
Mikhaylova M, Karpova A, Bär J, Bethge P, YuanXiang P, Chen Y, Zuschratter W, Behnisch T, Kreutz MR
Brain structure & function (2014) 2193: 843-60. . **ICC, IHC**

Selected General References

The vesicular GABA transporter, VGAT, localizes to synaptic vesicles in sets of glycinergic as well as GABAergic neurons.
Chaudhry FA et al. J. Neurosci. (1998) PubMed:9822734

Identification and characterization of the vesicular GABA transporter.
McIntire SL et al. Nature (1997) PubMed:9349821

Cloning of a functional vesicular GABA and glycine transporter by screening of genome databases.
Sagné C et al. FEBS Lett. (1997) PubMed:9395291

Uptake of GABA by rat brain synaptic vesicles isolated by a new procedure.
Hell JW et al. EMBO J. (1988) PubMed:2903047

Access the online factsheet including applicable protocols at https://sysy.com/product/131011C3 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.