

Rudolf-Wissell-Str. 28a 37079 Göttingen, Germany Phone: +49 551-50556-0 Fax: +49 551-50556-384 E-mail: sales@sysy.com Web: www.sysy.com

# GFP

Cat.No. 132 003C3; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

# Data Sheet

Reconstitution/ Storage	50 μg specific antibody, lyophilized. Affinity purified with the immunogen, fluorescence-labeled with Oyster 550. Albumin was added for stabilization. For <b>reconstitution</b> add 50 μl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Either add 1:1 (v/v) glycerol, then aliquot and store at -20°C until use, or store aliquots at -80°C without additives. Reconstitute immediately upon receipt! Avoid bright light when working with the antibody to minimize photo bleeching of the fluorescent dye.The mounting agent Aquatex <sup>®</sup> (Merck Chemicals) is not compatible with Oyster dyes! For detailed information, see back of the data sheet.
Applications	WB: N/A IP: N/A ICC: 1 : 500 IHC: 1 : 500 IHC-P: not tested yet
Label	Oyster 550
Immunogen	Recombinant protein corresponding to AA 1 to 238 from jellyfish GFP (UniProt Id: P42212)
Specificity	Recognizes GFP, mEGFP, superfolder GFP, most common CFP and YFP variants.Does not cross-react to mCherry, mRFP, dsRed, mTagBFP or their most common derivatives.

#### TO BE USED IN VITRO / FOR RESEARCH ONLY NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

## Background

**G**reen **f**luorescent **p**rotein **GFP** and its derivates have become universal tools in cell biology. These antibodies allow immunoprecipitation and visualization of GFP fusion proteins on immunoblots and by immunocytochemistry.

## **Selected General References**

Imaging into the future: visualizing gene expression and protein interactions with fluorescent proteins. van Roessel P et al. Nat. Cell Biol. (2002) PubMed:11780139

Illuminating the secretory pathway: when do we need vesicles? Stephens DJ et al. J. Cell. Sci. (2001) PubMed:11228150

Watching proteins in the wild: fluorescence methods to study protein dynamics in living cells. Chamberlain C et al. Traffic (2000) PubMed:11208065

Access the online factsheet including applicable protocols at <u>https://sysy.com/product/132003C3</u> or scan the QR-code.



# FAQ - How should I store my antibody?

### **Shipping Conditions**

• All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freezedried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- Unlabeled and biotin-labeled antibodies and control proteins should be stored at 4°C before reconstitution. They must not be stored in the freezer when still lyophilized! Temperatures below zero may cause loss of performance.
- Fluorescence-labeled antibodies should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

# Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 μl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## **Product Specific Hints for Storage**

#### Control proteins / peptides

• Store at -20°C to -80°C.

#### **Monoclonal Antibodies**

- Ascites and hybridoma supernatant should be stored at -20°C up to -80°C. Prolonged storage at 4°C is not recommended! Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### **Polyclonal Antibodies**

- **Crude antisera**: With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- Affinity purified antibodies: Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### **Fluorescence-labeled Antibodies**

• Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

# FAQ - How should I reconstitute my antibody?

## Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freezethaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.