

## VGLUT 1

Cat.No. 135 304; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/Storage	100 µl antiserum, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 5000 (AP staining) <b>IP:</b> yes <b>ICC:</b> 1 : 1000 up to 1 : 5000 <b>IHC:</b> 1 : 500 up to 1 : 1000 <b>IHC-P/FFPE:</b> 1 : 200 <b>EM:</b> yes <b>FACS:</b> yes
Immunogen	Recombinant protein corresponding to residues near the carboxy terminus of rat VGLUT 1 (UniProt Id: Q62634)
Reactivity	Reacts with: rat (Q62634), mouse (Q3TXX4), human (Q9P2U7), cow. Other species not tested yet.
Specificity	Specific for VGLUT 1. K.O.
Matching control	135-3P
Remarks	VGLUT 1 aggregates after boiling, making it necessary to run SDS-PAGE only with non-boiled samples.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Access the online factsheet including applicable protocols at <https://susy.com/product/135304> or scan the QR-code.



### Background

The vesicular **glutamate transporter 1 VGLUT 1**, also referred to as **BNPI** and **SLC17A7**, was originally identified as a brain specific phosphate transporter. Like the related VGLUT 2, VGLUT 1 is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane.

VGLUT 1 and VGLUT 2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

### Selected References for 135 304

- Quantitative comparison of glutamatergic and GABAergic synaptic vesicles unveils selectivity for few proteins including MAL2, a novel synaptic vesicle protein.  
Grønborg M, Pavlos NJ, Brunk I, Chua JJ, Münster-Wandowski A, Riedel D, Ahnert-Hilger G, Urlaub H, Jahn R  
The Journal of neuroscience : the official journal of the Society for Neuroscience (2010) 30(1): 2-12. . **ICC, IHC, EM**
- Prenatal interleukin 6 elevation increases glutamatergic synapse density and disrupts hippocampal connectivity in offspring.  
Mirabella F, Desiato G, Mancinelli S, Fossati G, Rasile M, Morini R, Markicevic M, Grimm C, Amegandjin C, Termini A, Peano C, et al.  
Immunity (2021) 54(11): 2611-2631.e8. . **WB, ICC, IHC; tested species: mouse**
- Inhibition of LRRK2 kinase activity promotes anterograde axonal transport and presynaptic targeting of α-synuclein.  
Brzozowski CF, Hijaz BA, Singh V, Gcwenza NZ, Kelly K, Boyden ES, West AB, Sarkar D, Volpicelli-Daley LA  
Acta neuropathologica communications (2021) 9(1): 180. . **WB, ICC, IHC; tested species: mouse**
- Extracellular transglutaminase-2, nude or associated with astrocytic extracellular vesicles, modulates neuronal calcium homeostasis.  
Tonoli E, Verduci I, Gabrielli M, Prada I, Forcaia G, Coveney C, Savoca MP, Boocock DJ, Sancini G, Mazzanti M, Verderio C, et al.  
Progress in neurobiology (2022) : 102313. . **WB, ICC; tested species: rat**
- Clathrin-independent endocytic retrieval of SV proteins mediated by the clathrin adaptor AP-2 at mammalian central synapses.  
López-Hernández T, Takenaka KI, Mori Y, Kongpracha P, Nagamori S, Hauke V, Takamori S  
eLife (2022) 11: . . **WB, ICC; tested species: mouse**
- Vesicular Glutamate Transporters (SLC17A6, 7, 8) Control Synaptic Phosphate Levels.  
Cheret C, Ganzella M, Preobraschenski J, Jahn R, Ahnert-Hilger G  
Cell reports (2021) 34(2): 108623. . **WB, ICC; tested species: human, mouse**
- SHANK2 Mutations Result in Dysregulation of the ERK1/2 Pathway in Human Induced Pluripotent Stem Cells-Derived Neurons and Shank2(-/-) Mice.  
Lutz AK, Pérez Arévalo A, Ioannidis V, Stirling N, Demestre M, Delorme R, Bourgeron T, Boeckers TM  
Frontiers in molecular neuroscience (2021) 14: 773571. . **WB, ICC; tested species: human**
- Chemico-genetic discovery of astrocytic control of inhibition in vivo.  
Takano T, Wallace JT, Baldwin KT, Purkey AM, Uezu A, Courtland JL, Soderblom EJ, Shimogori T, Maness PF, Eroglu C, Soderling SH, et al.  
Nature (2020) : . . **ICC, IHC; tested species: mouse**
- The small GTPase ARF6 regulates GABAergic synapse development.  
Kim H, Jung H, Jung H, Kwon SK, Ko J, Um JW  
Molecular brain (2020) 13(1): 2. . **ICC, IHC; tested species: rat**
- A CRHR1 antagonist prevents synaptic loss and memory deficits in a trauma-induced delirium-like syndrome.  
Cursano S, Battaglia CR, Urrutia-Ruiz C, Grabrucker S, Schön M, Bockmann J, Braumüller S, Radermacher P, Roselli F, Huber-Lang M, Boeckers TM, et al.  
Molecular psychiatry (2020) : . . **ICC, IHC; tested species: mouse**
- Acute knockdown of Depdc5 leads to synaptic defects in mTOR-related epileptogenesis.  
De Fusco A, Cerullo MS, Marte A, Michetti C, Romei A, Castroflorio E, Baulac S, Benfenati F  
Neurobiology of disease (2020) : 104822. . **ICC, IHC; tested species: mouse**
- Neurocan Contributes to Perineuronal Net Development.  
Schmidt S, Arendt T, Morawski M, Sonntag M  
Neuroscience (2020) 442: 69-86. . **WB, IHC; tested species: mouse**

# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.