SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

VGLUT1 (SLC17A7)

Cat.No. 135 318; Recombinant Guinea pig antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 1000 (AP staining) (see remarks) **IP**: not tested yet **ICC**: 1 : 500 **IHC**: 1 : 500 **IHC-P (FFPE)**: 1 : 4000
Clone	Gp68B7
Subtype	IgG2 (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the carboxy terminus of rat VGLUT 1. (UniProt Id: Q62634)
Reactivity	Reacts with: rat (Q62634), mouse (Q3TXX4), human (Q9P2U7). Other species not tested yet. Predicted to cross-react with cat due to high sequence homology.
Specificity	K.O. validated
Matching control	135-0P
Remarks	This antibody is a chimeric antibody based on the well-known monoclonal mouse antibody clone 68B7. The constant regions of the heavy and light chains have been replaced by Guinea pig specific sequences. Therefore, the antibody can be used with standard anti-Guinea pig secondary reagents. The antibody has been expressed in mammalian cells. **WB**: To avoid protein aggregation, do not heat samples for SDS-PAGE. This antibody is highly recommended for Western blot experiments.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The **v**esicular **glu**tamate **t**ransporter **1 VGLUT1**, also referred to as **BNPI** and **SLC17A7**, was originally identified as a brain specific phosphate transporter. Like the related VGLUT2, VGLUT1 is both necessary and sufficient for uptake and storage of glutamate and thus comprises the sole determinant for a glutamatergic phenotype. Both VGLUTs are different from the plasma membrane transporters in that they are driven by a proton electrochemical gradient across the vesicle membrane.
VGLUT1 and VGLUT2 show complementary expression patterns. Together, they are currently the best markers for glutamatergic nerve terminals and glutamatergic synapses.

Selected References for 135 318

T Cell-Mediated Targeting of Interneurons in Mice Shapes Hippocampal Remodeling and Epilepsy.
Galvis-Montes DS, Henning L, van Loo KMJ, Surges R, Schoch S, Becker AJ, Pitsch J
Annals of neurology (2026) : . . **IHC-P; tested species: mouse**

A postsynaptic GPR158-PLCXD2 complex controls spine apparatus abundance and dendritic spine maturation.
Verpoort B, Amado L, Vandensteen J, Leysen E, Dascenco D, Vandenbempt J, Lemmens I, Wauman J, Vennekens K, Escamilla-Ayala A, Freitas ACN, et al.
Developmental cell (2025) : . . **IHC; tested species: mouse**

Amygdala stress-responsive ensembles mediate distinct susceptibility to negative stress-induced remote depression in adolescent and adult mice.
You Y, Lv CT, Ni PT, Wang SJ, Liu L, Song YT, Hui Z, Huang Y, Liu Y, Tong K, Hao JR, et al.
Cell reports (2026) 454: 117281. . **IHC; tested species: mouse**

The complement C3-microglial axis in depression of Parkinson's disease: from mechanism to therapeutic intervention.
Yin Q, Ding M, Tang Y, Qi Y, Qin Y, Jin H, Li Y, Bao J, Ma S, Li Y, Ding H, et al.
EBioMedicine (2026) 129: 106325. . **IHC; tested species: mouse**

Synaptophysin density underestimates synapse counts in the human brain - An anatomical study on synaptoporin protein expression.
Woelfle S, Schön M, Pedro MT, Wagner J, Shiban E, Hong B, Tsoy Y, Boeckers TM
Neurobiology of disease (2026) 227: 107503. . **IHC; tested species: human**

Youth-associated protein TIMP2 regulates microglial state and function in healthy and aged mice.
Hemmer BM, Philippi SM, Ferreira AC, Petridis SF, Phan A, Castellano JM
Nature communications (2026) 171: . . **IHC; tested species: mouse**

Galectin-3 in microglia mediates neuroinflammation-induced cognitive dysfunction via selective elimination of excitatory synapses in hippocampal CA1.
Wu HP, Hu XY, Liu K, He QL, Zhu SY, Shi JY, Yang JJ, Fan D, Ji MH
Brain research (2026) 1887: 150382. . **IHC; tested species: mouse**

Selected General References

Identification of a vesicular glutamate transporter that defines a glutamatergic phenotype in neurons.
Takamori S et al. Nature (2000) PubMed:11001057

Uptake of glutamate into synaptic vesicles by an inorganic phosphate transporter.
Bellocchio EE et al. Science (2000) PubMed:10938000

The localization of the brain-specific inorganic phosphate transporter suggests a specific presynaptic role in glutamatergic transmission.
Bellocchio EE et al. J. Neurosci. (1998) PubMed:9786972

Access the online factsheet including applicable protocols at https://sysy.com/product/135318 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.