

ERC1b/2

Cat.No. 143-0P; control peptide, 100 µg peptide (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg peptide, lyophilized. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Control peptides should be stored at -20°C when still lyophilized! For detailed information, see back of the data sheet.
Immunogen	Synthetic peptide corresponding to AA 939 to 948 from rat Erc1b (UniProt Id: Q811U3-1)
Recommended dilution	Optimal concentrations should be determined by the end-user.
Matching antibodies	143 003, 143 004
Remarks	This control peptide consists of the synthetic peptide CDQDEEEGIWA (aa 939 - 948 in rat ERC 1b) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of protein needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

ELKS, also referred to as **ERCs** (ERC 1 and ERC 2) and **CAST**, are related proteins which share an identical C-terminal sequence. They interact with the conserved RIM PDZ domain via an unusual PDZ binding motif. Two splice variants of ERC 1 (1a and 1b) have been described. ERC 1b (CAST 2a) binds to RIM and is expressed exclusively in the brain. ERC 1a is a ubiquitously expressed cytosolic protein. ERC 2 (CAST 1) is only expressed as a single RIM binding variant.

All ERCs have been shown to interact with Rab 6, a protein involved in membrane trafficking at the Golgi complex. The function of these proteins has not been determined yet. They may link Rab 6 mediated non-neuronal membrane traffic at the Golgi complex to neuronal membrane traffic at the active zone executed via RIMs.

Selected General References

Physical and functional interaction of the active zone proteins, CAST, RIM1, and Bassoon, in neurotransmitter release. Takao-Rikitsu E et al. J. Cell Biol. (2004) PubMed:14734538

CAST2: identification and characterization of a protein structurally related to the presynaptic cytomatrix protein CAST. Deguchi-Tawarada M et al. Genes Cells (2004) PubMed:14723704

Interaction of the ERC family of RIM-binding proteins with the liprin-alpha family of multidomain proteins. Ko J et al. J. Biol. Chem. (2003) PubMed:12923177

Cast: a novel protein of the cytomatrix at the active zone of synapses that forms a ternary complex with RIM1 and munc13-1. Ohtsuka T et al. J. Cell Biol. (2002) PubMed:12163476

A family of RIM-binding proteins regulated by alternative splicing: Implications for the genesis of synaptic active zones. Wang Y et al. Proc. Natl. Acad. Sci. U.S.A. (2002) PubMed:12391317

Access the online factsheet including applicable protocols at <https://sysy.com/product/143-0P> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.