

## Arc

Cat.No. 156 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Azide was added before lyophilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> not recommended <b>IP:</b> yes <b>ICC:</b> not tested yet <b>IHC:</b> not tested yet <b>IHC-P:</b> not tested yet <b>ELISA:</b> yes (see remarks)
Clone	48B2
Subtype	IgG2b (κ light chain)
Immunogen	Full-length recombinant mouse Arc (UniProt Id: Q9WV31)
Epitop	AA 1 to 214 from mouse Arc (UniProt Id: Q9WV31)
Reactivity	Reacts with: rat (Q63053), mouse (Q9WV31). Other species not tested yet.
Specificity	Specific for Arc.
Remarks	<b>ELISA:</b> Suitable as capture antibody for sandwich-ELISA with cat. no. <a href="#">156 003</a> as detector antibody. The ELISA-protocol for membrane proteins is recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Immediate-early genes (IEGs) are rapidly induced after patterned synaptic activity. Genes that are involved in this complex response code for transcription and growth factors, metabolic and signaling enzymes, small GTP binding proteins and structural proteins. Some of these proteins may play a crucial role in long term plasticity which is important for learning processes. The activity regulated cytoskeleton associated protein **Arc** or **Arg 3.1** is enriched in dendrites and colocalizes with F-Actin. Direct interaction of Arc with actin has also been demonstrated by biochemical studies.

### Selected References for 156 011

Synapsin-dependent reserve pool of synaptic vesicles supports replenishment of the readily releasable pool under intense synaptic transmission.  
Vasileva M, Horstmann H, Geumann C, Gitler D, Kuner T  
The European journal of neuroscience (2012) 368: 3005-20. . **ELISA**

### Selected General References

Regulation of activity-regulated cytoskeleton protein (Arc) mRNA after acute and chronic electroconvulsive stimulation in the rat.  
Larsen MH et al. Brain Res. (2005) PubMed:16309632

Memory-influencing intra-basolateral amygdala drug infusions modulate expression of Arc protein in the hippocampus.  
McIntyre CK et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PubMed:16020527

Sparse, environmentally selective expression of Arc RNA in the upper blade of the rodent fascia dentata by brief spatial experience.  
Chawla MK et al. Hippocampus (2005) PubMed:15920719

Experience-dependent coincident expression of the effector immediate-early genes arc and Homer 1a in hippocampal and neocortical neuronal networks.  
Vazdarjanova A et al. J. Neurosci. (2002) PubMed:12451105

Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory.  
Guzowski JF et al. J. Neurosci. (2000) PubMed:10818134

The activity-regulated cytoskeletal-associated protein arc is expressed in different striosome-matrix patterns following exposure to amphetamine and cocaine.  
Tan A et al. J. Neurochem. (2000) PubMed:10800951

Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles.  
Guzowski JF et al. Nat. Neurosci. (1999) PubMed:10570490

Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites.  
Steward O et al. Neuron (1998) PubMed:9808461

Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites.  
Lyford GL et al. Neuron (1995) PubMed:7857651

Access the online factsheet including applicable protocols at <https://sysy.com/product/156011> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.