SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

# Arc

**Cat.No. 156 111; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)**

## Data Sheet

| | |
|---|---|
| Reconstitution/ Storage | 100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization.<br>For **reconstitution** add 100 µl $H_2O$ to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use.<br>Antibodies should be stored at +4°C when still lyophilized. Do not freeze!<br>For detailed information, see back of the data sheet. |
| Applications | **WB**: 1 : 1000 (AP staining)<br>**IP**: not tested yet<br>**ICC**: not tested yet<br>**IHC**: 1 : 500<br>**IHC-P (FFPE)**: not tested yet |
| Clone | 73A5 |
| Subtype | IgG1 (κ light chain) |
| Immunogen | Synthetic peptide corresponding to residues near the carboxy terminus of mouse Arc (UniProt Id: Q9WV31) |
| Reactivity | Reacts with: mouse (Q9WV31), rat (Q63053).<br>Other species not tested yet. |
| Specificity | Specific for Arc. |

**TO BE USED IN VITRO / FOR RESEARCH ONLY**
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Immediate-early genes (IEGs) are rapidly induced after patterned synaptic activity. Genes that are involved in this complex response code for transcription and growth factors, metabolic and signaling enzymes, small GTP binding proteins and structural proteins. Some of these proteins may play a crucial role in long term plasticity which is important for learning processes.
The **a**ctivity **r**egulated **c**ytoskeleton associated protein **Arc** or **Arg 3.1** is enriched in dendrites and colocalizes with F-Actin. Direct interaction of Arc with actin has also been demonstrated by biochemical studies.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

## Selected General References

Regulation of activity-regulated cytoskeleton protein (Arc) mRNA after acute and chronic electroconvulsive stimulation in the rat.
Larsen MH et al. Brain Res. (2005) PubMed:16309632

Memory-influencing intra-basolateral amygdala drug infusions modulate expression of Arc protein in the hippocampus.
McIntyre CK et al. Proc. Natl. Acad. Sci. U.S.A. (2005) PubMed:16020527

Sparse, environmentally selective expression of Arc RNA in the upper blade of the rodent fascia dentata by brief spatial experience.
Chawla MK et al. Hippocampus (2005) PubMed:15920719

Experience-dependent coincident expression of the effector immediate-early genes arc and Homer 1a in hippocampal and neocortical neuronal networks.
Vazdarjanova A et al. J. Neurosci. (2002) PubMed:12451105

Inhibition of activity-dependent arc protein expression in the rat hippocampus impairs the maintenance of long-term potentiation and the consolidation of long-term memory.
Guzowski JF et al. J. Neurosci. (2000) PubMed:10818134

The activity-regulated cytoskeletal-associated protein arc is expressed in different striosome-matrix patterns following exposure to amphetamine and cocaine.
Tan A et al. J. Neurochem. (2000) PubMed:10800951

Environment-specific expression of the immediate-early gene Arc in hippocampal neuronal ensembles.
Guzowski JF et al. Nat. Neurosci. (1999) PubMed:10570490

Synaptic activation causes the mRNA for the IEG Arc to localize selectively near activated postsynaptic sites on dendrites.
Steward O et al. Neuron (1998) PubMed:9808461

Arc, a growth factor and activity-regulated gene, encodes a novel cytoskeleton-associated protein that is enriched in neuronal dendrites.
Lyford GL et al. Neuron (1995) PubMed:7857651

Access the online factsheet including applicable protocols at https://sysy.com/product/156111 or scan the QR-code.

# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

# FAQ - How should I reconstitute my antibody?

## Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.