

## Vimentin

Cat.No. 172 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O, then aliquot and store at -20°C until use. For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 <b>IHC:</b> 1 : 500 <b>IHC_P:</b> 1 : 250 up to 1 : 1000
Immunogen	Recombinant protein corresponding to AA 1 to 466 from mouse Vimentin (UniProt Id: P20152)
Reactivity	Reacts with: mouse (P20152), rat (P31000). Other species not tested yet.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Vimentin** belongs to the family of intermediate filaments that can be subdivided into six major groups based on sequence similarity. Vimentin belongs to the type III category and is the predominant subunit protein of intermediate filaments in tissues of mesenchymal origin. Like other intermediate filaments it plays a role in the cytoskeletal organization and maintenance of cell shape and morphology.

## Selected General References

Architecture of the vimentin cytoskeleton is modified by perturbation of the GTPase ARF1.  
Styers ML, Kowalczyk AP, Faundez V  
Journal of cell science (2006) 119Pt 17: 3643-54. .

A direct interaction between actin and vimentin filaments mediated by the tail domain of vimentin.  
Esue O, Carson AA, Tseng Y, Wirtz D  
The Journal of biological chemistry (2006) 28141: 30393-9. .

Ultrastructure of intermediate filaments of nestin- and vimentin-immunoreactive astrocytes in organotypic slice cultures of hippocampus.  
Miyaguchi K  
Journal of structural biology (1997) 1201: 61-8. .

The cytoskeleton of primary astrocytes in culture contains actin, glial fibrillary acidic protein, and the fibroblast-type filament protein, vimentin.  
Chiu FC, Norton WT, Fields KL  
Journal of neurochemistry (1981) 371: 147-55. .

Vimentin: a phosphoprotein under hormonal regulation.  
Browning ET, Sanders MM  
The Journal of cell biology (1981) 903: 803-8. .

The synthesis and distribution of desmin and vimentin during myogenesis in vitro.  
Gard DL, Lazarides E  
Cell (1980) 191: 263-75. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/172004> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.