SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

# GFAP

**Cat.No. 173 008; Recombinant rabbit antibody, 50 µg recombinant IgG (lyophilized)**

## Data Sheet

| | |
|---|---|
| Reconstitution/ Storage | 50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For **reconstitution** add 50 µl $H_2O$ to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use.<br>Antibodies should be stored at +4°C when still lyophilized. Do not freeze!<br>For detailed information, see back of the data sheet. |
| Applications | **WB**: 1 : 1000 (AP staining) (see remarks)<br>**IP**: not tested yet<br>**ICC**: 1 : 1000<br>**IHC**: 1 : 2000<br>**IHC-P (FFPE)**: 1 : 5000 |
| Clone | Rb134B1 |
| Subtype | IgG1 (κ light chain) |
| Immunogen | full-length recombinant human GFAP (UniProt Id: P14136) |
| Epitop | AA 391 to 405 from human GFAP (UniProt Id: P14136) |
| Reactivity | Reacts with: human (P14136), rat (P47819), mouse (P03995), cow.<br>No signal: zebrafish.<br>Other species not tested yet. |
| Specificity | Specific for GFAP isoform 1. K.O. validated |
| Matching control | 173-0P |
| Remarks | This antibody is a chimeric antibody based on the well known monoclonal mouse antibody clone 134B1. The constant regions of the heavy and light chains have been replaced by rabbit specific sequences. Therefore, the antibody can be used with standard anti-rabbit secondary reagents. The antibody has been expressed in mammalian cells.<br>**WB**: In Western blots, the monoclonal GFAP antibodies are less sensitive than the polyclonal rabbit antibody. |

**TO BE USED IN VITRO / FOR RESEARCH ONLY**
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**G**lial **f**ibrillary **a**cidic **p**rotein **GFAP** is a glial-specific member of the intermediate filament protein family. This group comprises cell type-specific filamentous proteins with similar structure and function as scaffold for cytoskeleton assembly and maintenance.
Frequently, neural stem cells also express GFAP. In addition many types of brain tumors, probably derived from astrocytic cells, heavily express GFAP. This protein is also found in the lens epithelium, Kupffer cells of the liver, in some cells in salivary tumors and others.
Point-mutations in the GFAP gene have been correlated to Alexander disease, a fatal leukoencephalopathy that leads to the dysmyelination or demyelination of the central nervous system.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

## Selected References for 173 008

Mitochondrial Ca2+ efflux controls neuronal metabolism and long-term memory across species.
Amrapali Vishwanath A, Comyn T, Mira RG, Brossier C, Pascual-Caro C, Faour M, Boumendil K, Chintaluri C, Ramon-Duaso C, Fan R, Ghosh K, et al.
Nature metabolism (2026) 82: 467-488. . **ICC; tested species: rat**

Cortical organoid-derived models of the melanoma brain metastatic niche enable prioritization of cancer-targeting drugs.
Krieg K, Materna-Reichelt S, Naber T, Rachad FZ, Kauven P, Weller A, Haferkamp U, Wittich A, Zaliani A, Woo MS, Walkenhorst M, et al.
Cell reports methods (2025) : 101236. . **IHC-P; tested species: human**

## Selected General References

Loss of glial fibrillary acidic protein (GFAP) impairs Schwann cell proliferation and delays nerve regeneration after damage.
Triolo D et al. J. Cell. Sci. (2006) PubMed:16988027

Asymptomatic hereditary Alexander's disease caused by a novel mutation in GFAP.
Shiihara T et al. J. Neurol. Sci. (2004) PubMed:15465095

Glial fibrillary acidic protein: GFAP-thirty-one years (1969-2000).
Eng LF et al. Neurochem. Res. (2000) PubMed:11059815

GFAP-positive and myelin marker-positive glia in normal and pathologic environments.
Dyer CA et al. J. Neurosci. Res. (2000) PubMed:10797544

Expression of GFAP immunoreactivity during development of long fiber tracts in the rat CNS.
Valentino KL et al. Brain Res. (1983) PubMed:6627026

Glial fibrillary acidic protein (GFAP): purification from human fibrillary astrocytoma, development and validation of a radioimmunoassay for GFAP-like immunoactivity.
Palfreyman JW et al. J. Neurol. Sci. (1979) PubMed:438840

Determination of glial fibrillary acidic protein (GFAP) in human brain tumors.
Jacque CM et al. J. Neurol. Sci. (1978) PubMed:624958

Access the online factsheet including applicable protocols at https://sysy.com/product/173008 or scan the QR-code.

# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

# FAQ - How should I reconstitute my antibody?

## Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.