

GluA1 (AMPA1)

Cat.No. 182 511; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 500 up to 1 : 1000 (AP-staining) IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 up to 1 : 2000 IHC-P (FFPE): 1 : 100
Clone	SY-70D11
Subtype	IgG2b (κ light chain)
Immunogen	Synthetic peptide corresponding to C-terminal residues of rat GluA1 (UniProt Id: P19490)
Reactivity	Reacts with: mouse (P23818), rat (P19490). Other species not tested yet.
Matching control	182-01P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Ionotropic **glutamate receptors (iGluRs)** mediate rapid excitatory neurotransmission in the mammalian CNS. They can be subdivided into three major groups, the **AMPA/GluA**, NMDA/GluN and kainate/GluK receptors (KAR). mRNAs coding for glutamate receptors are substrates for an adenosine deaminase acting on RNA (ADAR) that increases the diversity of these proteins. Glutamate receptors of the AMPA subtype are monovalent cation channels and are composed of the four AMPA subunits **GluA1**, GluA2, GluA3, and GluA4.

Selected General References

- A nomenclature for ligand-gated ion channels.
Collingridge GL et al. Neuropharmacology (2009) PubMed:18655795
- Differential regulation of dendrite complexity by AMPA receptor subunits GluR1 and GluR2 in motor neurons.
Priithviraj R et al. Dev Neurobiol (2008) PubMed:18000827
- Influence of environmental enrichment on steady-state mRNA levels for EAAC1, AMPA1 and NMDA2A receptor subunits in rat hippocampus.
Andin J et al. Brain Res. (2007) PubMed:17854777
- Identification and characterization of a novel phosphorylation site on the GluR1 subunit of AMPA receptors.
Lee HK et al. Mol. Cell. Neurosci. (2007) PubMed:17689977
- Receptor occupancy and channel-opening kinetics: a study of GLUR1 L497Y AMPA receptor.
Pei W et al. J. Biol. Chem. (2007) PubMed:17545169
- Differential localization of the GluR1 and GluR2 subunits of the AMPA-type glutamate receptor among striatal neuron types in rats.
Deng YP et al. J. Chem. Neuroanat. (2007) PubMed:17446041
- Widespread expression of the AMPA receptor GluR2 subunit at glutamatergic synapses in the rat spinal cord and phosphorylation of GluR1 in response to noxious stimulation revealed with an antigen-unmasking method.
Nagy GG et al. J. Neurosci. (2004) PubMed:15215299
- Phosphorylation of the AMPA receptor GluR1 subunit is required for synaptic plasticity and retention of spatial memory.
Lee HK et al. Cell (2003) PubMed:12628184
- NMDA induces long-term synaptic depression and dephosphorylation of the GluR1 subunit of AMPA receptors in hippocampus.
Lee HK et al. Neuron (1998) PubMed:9856470
- Differential expression of glutamate receptor genes (GluR1-5) in the rat retina.
Hughes TE et al. Vis. Neurosci. (1992) PubMed:1310870
- The characterization and localization of the glutamate receptor subunit GluR1 in the rat brain.
Rogers SW et al. J. Neurosci. (1991) PubMed:1652625

Access the online factsheet including applicable protocols at <https://sysy.com/product/182511> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.