

## PACSIN1

Cat.No. 196 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 up to 1 : 1000 <b>IHC:</b> not tested yet <b>IHC_P:</b> not tested yet
Immunogen	Recombinant protein corresponding to AA 293 to 384 from mouse PACSIN1 (UniProt Id: Q61644)
Reactivity	Reacts with: rat (Q9Z0W5), mouse (Q61644), human (Q9BY11), zebrafish. Other species not tested yet.
Specificity	K.D. PubMed: <a href="https://pubmed.ncbi.nlm.nih.gov/30683896/">30683896</a>
Matching control	196-0P

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

This protein has been described as **PACSIN 1** (for **PKC** and **CK2** substrate in neurons) and **syndapin 1** (**synaptic, dynamin-associated protein I**). It is a multidomain phospho-protein that interacts via its SH3 domain with dynamin 1, synaptojanin, synapsin and N-WASP. Since N-WASP is an activator of the Arp2/3 complex, syndapin may link membrane trafficking and the actin cytoskeleton. Up to now, 3 isoforms have been described. Syndapin 1 shows a neuronal, syndapin 3 a muscular and syndapin 2 an ubiquitous expression profile.

### Selected References for 196 003

Investigation of F-BAR domain PACSIN proteins uncovers membrane tubulation function in cilia assembly and transport. Insinna C, Lu Q, Teixeira I, Harned A, Semler EM, Stauffer J, Magidson V, Tiwari A, Kenworthy AK, Narayan K, Westlake CJ, et al. Nature communications (2019) 101: 428. . **WB, IHC; KD verified; tested species: human, zebrafish**

### Selected General References

Syndapin I is the phosphorylation-regulated dynamin I partner in synaptic vesicle endocytosis. Anggono V, Smillie KJ, Graham ME, Valova VA, Cousin MA, Robinson PJ. Nature neuroscience (2006) 96: 752-60. .

Syndapin oligomers interconnect the machineries for endocytic vesicle formation and actin polymerization. Kessels MM, Qualmann B. The Journal of biological chemistry (2006) 28119: 13285-99. .

The syndapin protein family: linking membrane trafficking with the cytoskeleton. Kessels MM, Qualmann B. Journal of cell science (2004) 117Pt 15: 3077-86. .

All three PACSIN isoforms bind to endocytic proteins and inhibit endocytosis. Modregger J, Ritter B, Witter B, Paulsson M, Plomann M. Journal of cell science (2000) 113 Pt 24: 4511-21. .

Syndapin isoforms participate in receptor-mediated endocytosis and actin organization. Qualmann B, Kelly RB. The Journal of cell biology (2000) 1485: 1047-62. .

Syndapin I, a synaptic dynamin-binding protein that associates with the neural Wiskott-Aldrich syndrome protein. Qualmann B, Roos J, DiGregorio PJ, Kelly RB. Molecular biology of the cell (1999) 102: 501-13. .

PACSIN, a brain protein that is upregulated upon differentiation into neuronal cells. Plomann M, Lange R, Vopper G, Cremer H, Heinlein UA, Scheff S, Baldwin SA, Leitges M, Cramer M, Paulsson M, Barthels D, et al. European journal of biochemistry (1998) 2561: 201-11. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/196003> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.