

## Calbindin D28k

Cat.No. 214 011C3; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized, fluorescence-labeled with Oyster 550. Albumin was added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Either add 1:1 (v/v) glycerol, then aliquot and store at -20°C until use, or store aliquots at -80°C without additives. Reconstitute immediately upon receipt! Avoid bright light when working with the antibody to minimize photo bleaching of the fluorescent dye. The mounting agent Aquatex® (Merck Chemicals) is not compatible with Oyster dyes! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> N/A <b>IP:</b> N/A <b>ICC:</b> not tested yet <b>IHC:</b> 1 : 200 <b>IHC_P:</b> not tested yet
Label	Oyster 550
Clone	351C10
Subtype	IgG1 (κ light chain)
Immunogen	Recombinant protein corresponding to AA 3 to 251 from human CalbindinD28k (UniProt Id: P05937)
Reactivity	Reacts with: human (P05937), rat (P07171), mouse (P12658), zebrafish. Other species not tested yet.
Matching control	214-0P

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Access the online factsheet including applicable protocols at <https://sysy.com/product/214011C3> or scan the QR-code.



### Background

Two isoforms of the vitamin D-dependent Ca-binding proteins have been described so far: **calbindin D28k**, also referred to as CALB 1, D-28k, and CAB 27, and calbindin D29k, also known as calretinin. These proteins are expressed in cells that have to handle a high calcium influx such as brain, bone, teeth, inner ear and others. Calbindins are believed to regulate cellular activity by suppressing or buffering intracellular calcium. In the brain calbindin D28k is a useful marker for specific neuronal cell types. It is particularly concentrated in the dendrites and perikarya of cerebellar Purkinje cells, but is also found in many GABAergic interneurons in the cortex.

### Selected References for 214 011C3

Stem cell regionalization during olfactory bulb neurogenesis depends on regulatory interactions between Vax1 and Pax6.  
Coré N, Erni A, Hoffmann HM, Mellon PL, Saurin AJ, Beclin C, Cremer H  
eLife (2020) 9: . . . **IHC; tested species: mouse**

AMIGO1 promotes axon growth and territory matching in the retina.

Soto F, Shen N, Kerschensteiner D

The Journal of neuroscience : the official journal of the Society for Neuroscience (2022) : . . . **IHC; tested species: mouse**

### Selected General References

Influence of the "open field" exposure on calbindin D28K, calretinin, and parvalbumin containing cells in the rat midbrain - developmental study.

Klejbor I, Ludkiewicz B, Domaradzka-Pytel B, Spodnik JH, Dziwiatkowski J, Morys J

Journal of physiology and pharmacology : an official journal of the Polish Physiological Society (2006) 571: 149-64. .

Calbindin D-28 and microtubule-associated protein-2: their use as sensitive immunohistochemical markers of cerebellar neurotoxicity in a regulatory toxicity study.

Haworth R, McCormack N, Selway S, Pilling AM, Williams TC

Experimental and toxicologic pathology : official journal of the Gesellschaft für Toxikologische Pathologie (2006) 575-6: 419-26. .

Mutational analysis of dendritic Ca<sup>2+</sup> kinetics in rodent Purkinje cells: role of parvalbumin and calbindin D28k.

Schmidt H, Stiefel KM, Racay P, Schwaller B, Eilers J

The Journal of physiology (2003) 551Pt 1: 13-32. .

Calbindin in cerebellar Purkinje cells is a critical determinant of the precision of motor coordination.

Barski JJ, Hartmann J, Rose CR, Hoebeek F, Mörl K, Noll-Hussong M, De Zeeuw CI, Konnerth A, Meyer M

The Journal of neuroscience : the official journal of the Society for Neuroscience (2003) 238: 3469-77. .

'New' functions for 'old' proteins: the role of the calcium-binding proteins calbindin D-28k, calretinin and parvalbumin, in cerebellar physiology. Studies with knockout mice.

Schwaller B, Meyer M, Schiffmann S

Cerebellum (London, England) (2002) 14: 241-58. .

Synthesis of calbindin-D28K during mineralization in human bone marrow stromal cells.

Fauchoux C, Bareille R, Amedee J

The Biochemical journal (1998) 333 ( Pt 3): 817-23. .

Calbindin-D in peripheral nerve cells is vitamin D and calcium dependent.

Lee YS, Taylor AN, Reimers TJ, Edelstein S, Fullmer CS, Wasserman RH

Proceedings of the National Academy of Sciences of the United States of America (1987) 8420: 7344-8. .

# FAQ - How should I store my antibody?

## Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C.

### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.