

## c-Fos

Cat.No. 226 017; Monoclonal rat antibody, 100 µg purified IgG (lyophilized)

## Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 up to 1 : 1000 <b>IHC:</b> 1 : 1000 up to 1 : 5000 (see remarks) <b>IHC_P:</b> 1 : 100 up to 1 : 200 <b>FACS:</b> yes
Clone	108B5H5
Subtype	IgG2a (κ light chain)
Immunogen	Synthetic peptide corresponding to residues near the amino terminus of rat c-Fos (UniProt Id: P12841)
Reactivity	Reacts with: mouse (P01101), rat (P12841), human (P01100). Other species not tested yet.
Matching control	226-0P
Remarks	<b>IHC:</b> For best results tissue sections should be stored at -20°C in cryoprotectant solution. Prolonged storage at 4°C leads to a substantial loss of signal.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

Access the online factsheet including applicable protocols  
at <https://sysy.com/product/226017> or scan the QR-code.



## Background

The Fos gene family consists of 4 members: **c-Fos**, FosB, FosL1, and FosL2, also called Fos related antigen 1 and 2 (FRA1 and 2). These leucine zipper proteins can dimerize with proteins of the JUN family leading to the formation of the transcription factor complex AP1 (1).  
The expression of Fos proteins is rapidly and transiently induced by different extracellular stimuli such as growth factors, cytokines, neurotransmitters, polypeptide hormones, stress (2).  
In addition Fos proteins can be phosphorylated by ERK kinases modulating transcriptional activity, protein stability and localization (3). c-Fos is the homologue to the Finkel-Biskis-Jenkins (FBJ) murine osteosarcoma virus oncogene (4).

## Selected References for 226 017

- Transcriptional dynamics of murine motor neuron maturation in vivo and in vitro.  
Patel T, Hammelman J, Aziz S, Jang S, Closser M, Michaels TL, Blum JA, Gifford DK, Wichterle H  
Nature communications (2022) 131: 5427. . **IHC, ICC; tested species: mouse**
- Adrenergic signalling to astrocytes in anterior cingulate cortex contributes to pain-related aversive memory in rats.  
Iqbal Z, Lei Z, Ramkrishnan AS, Liu S, Hasan M, Akter M, Lam YY, Li Y  
Communications biology (2023) 61: 10. . **IHC; tested species: rat**
- Cry1Δ11 mutation induces ADHD-like symptoms through hyperactive dopamine D1 receptor signaling.  
Liu D, Xie Z, Gu P, Li X, Zhang Y, Wang X, Chen Z, Deng S, Shu Y, Li JD  
JCI insight (2023) 816: . . **IHC; tested species: mouse**
- Chronic Pb Exposure Induces Anxiety and Depression-like Behaviors in Mice via Excitatory Neuronal Hyperexcitability in Ventral Hippocampal Dentate Gyrus.  
Zhou R, Huang C, Bi N, Li L, Li C, Gu X, Song Y, Wang HL  
Environmental science & technology (2023) 5733: 12222-12233. . **IHC; tested species: mouse**
- Bidirectional Regulation of Motor Circuits Using Magnetogenetic Gene Therapy.  
Unda SR, Marongiu R, Pomeranz LE, Dyke JP, Fung EK, Grosenick L, Zirkel R, Antoniazzi AM, Norman S, Liston CM, Schaffer C, et al.  
bioRxiv : the preprint server for biology (2023) : . . **IHC; tested species: mouse**
- ΔFosB accumulation in hippocampal granule cells drives cFos pattern separation during spatial learning.  
Lamothe-Molina PJ, Franzelin A, Beck L, Li D, Auksutat L, Fieblinger T, Laprell L, Alhbeck J, Gee CE, Kneussel M, Engel AK, et al.  
Nature communications (2022) 131: 6376. . **IHC; tested species: mouse**
- Thalamo-hippocampal pathway regulates incidental memory capacity in mice.  
Torromino G, Loffredo V, Cavezza D, Sonsini G, Esposito F, Crevenna AH, Gioffrè M, De Risi M, Treves A, Griguoli M, De Leonibus E, et al.  
Nature communications (2022) 131: 4194. . **IHC; tested species: mouse**
- Reprogramming reactive glia into interneurons reduces chronic seizure activity in a mouse model of mesial temporal lobe epilepsy.  
Lentini C, d'Orange M, Marichal N, Trottmann MM, Vignoles R, Foucault L, Verrier C, Massera C, Raineteau O, Conzelmann KK, Rival-Gervier S, et al.  
Cell stem cell (2021) : . . **IHC; tested species: mouse**
- Non-invasive optogenetics with ultrasound-mediated gene delivery and red-light excitation.  
Pouliopoulos AN, Murillo MF, Noel RL, Batts AJ, Ji R, Kwon N, Yu H, Tong CK, Gelinas JN, Araghy DK, Hussaini SA, et al.  
Brain stimulation () 154: 927-941. . **IHC; tested species: mouse**

## Selected General References

- Existence of different Fos/Jun complexes during the G0-to-G1 transition and during exponential growth in mouse fibroblasts: differential role of Fos proteins.  
Kovary K, Bravo R  
Molecular and cellular biology (1992) 1211: 5015-23. .
- Regulation of the transcriptional activity of c-Fos by ERK. A novel role for the prolyl isomerase PIN1.  
Monje P, Hernández-Losa J, Lyons RJ, Castellone MD, Gutkind JS  
The Journal of biological chemistry (2005) 28042: 35081-4. .
- Oncogenes in rheumatoid arthritis.  
Müller-Ladner U, Kriegsmann J, Gay RE, Gay S  
Rheumatic diseases clinics of North America (1995) 213: 675-90. .

## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.