

Claudin11

Cat.No. 241 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) (see remarks) IP: not tested yet ICC: 1 : 500 IHC: 1 : 500 (see remarks) IHC-P (FFPE): 1 : 200 up to 1 : 500
Immunogen	Synthetic peptide corresponding to AA 188 to 207 from rat Claudin11 (UniProt Id: Q99P82)
Reactivity	Reacts with: human (O75508), rat (Q99P82), mouse (Q60771), cow, pig. Other species not tested yet.
Matching control	241-0P
Remarks	WB: To avoid protein aggregation, do not heat samples for SDS-PAGE. IHC: Antigen retrieval with citrate buffer pH 6 can be applied to improve the signal to noise ratio.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Claudin 11, also known as **Oligodendrocyte specific protein (OSP)**, is one of the most abundant components of CNS myelin where it locates to the interlamellar strands in myelin sheaths. To date more than 20 members of the claudin family have been identified. They are expressed in epithelial and endothelial cells in all tight junction bearing tissues. Claudins share a common transmembrane topology with 4 transmembrane spanning domains and two extracellular loops.

Selected References for 241 003

Immunosignals of Oligodendrocyte Markers and Myelin-Associated Proteins Are Critically Affected after Experimental Stroke in Wild-Type and Alzheimer Modeling Mice of Different Ages.
Michalski D, Keck AL, Grosche J, Martens H, Härtig W
Frontiers in cellular neuroscience (2018) 12: 23. . **IHC; tested species: mouse**

Selected General References

Tight junctions potentiate the insulative properties of small CNS myelinated axons.
Devaux J et al. J. Cell Biol. (2008) PubMed:19047465

Compartmentalization established by claudin-11-based tight junctions in stria vascularis is required for hearing through generation of endocochlear potential.
Kitajiri S et al. J. Cell. Sci. (2004) PubMed:15456848

OSP/claudin-11 forms a complex with a novel member of the tetraspanin super family and beta1 integrin and regulates proliferation and migration of oligodendrocytes.
Tiwari-Woodruff SK et al. J. Cell Biol. (2001) PubMed:11309411

Developmental expression of OSP/claudin-11.
Bronstein JM et al. J. Neurosci. Res. (2000) PubMed:10797530

Claudin-11/OSP-based tight junctions of myelin sheaths in brain and Sertoli cells in testis.
Morita K et al. J. Cell Biol. (1999) PubMed:10225958

Oligodendrocyte-specific protein (OSP) is a major component of CNS myelin.
Bronstein JM et al. J. Neurosci. Res. (1997) PubMed:9418959

Access the online factsheet including applicable protocols at <https://sysy.com/product/241003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.