

Cre-Recombinase

Cat.No. 257 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: not tested yet IP: not tested yet ICC: 1 : 500 up to 1 : 1000 IHC: 1 : 500 IHC_P: 1 : 500
Immunogen	Full length recombinant Cre-recombinase from Bacteriophage P1 (UniProt Id: P06956)
Specificity	Recognizes Cre- and iCre-recombinase.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Cre-Recombinase is a 38 kDa type 1 topoisomerase from bacteriophage P1. It is highly specific for a 34 bp DNA sequence (loxP) found in P1 DNA and catalyzes site-specific recombination between two 34-base-pair LOXP recognition sites. Its role is to maintain the phage genome as a monomeric unit-copy plasmid in the lysogenic state. Its highly specific catalytic activity makes it a valuable tool for the generation of conditional or tissue specific mutants.

Selected References for 257 004

Ultrastructural Imaging of Activity-Dependent Synaptic Membrane-Trafficking Events in Cultured Brain Slices.
Imig C, López-Murcia FJ, Maus L, García-Plaza IH, Mortensen LS, Schwark M, Schwarze V, Angibaud J, Nägerl UV, Taschenberger H, Brose N, et al.
Neuron (2020) : . . **IHC; tested species: mouse**

Endogenous opioid system modulates conditioned cocaine reward in a sex-dependent manner.
Matsumura K, Nicot A, Choi IB, Asokan M, Le NN, Natividad LA, Dobbs LK
Addiction biology (2023) 2810: e13328. . **IHC; tested species: mouse**

Striatonigrostriatal circuit architecture for disinhibition of dopamine signaling.
Ambrosi P, Lerner TN
Cell reports (2022) 407: 111228. . **IHC; tested species: mouse**

Cone-Driven Retinal Responses Are Shaped by Rod But Not Cone HCN1.
Lankford CK, Umino Y, Poria D, Kefalov V, Solessio E, Baker SA
The Journal of neuroscience : the official journal of the Society for Neuroscience (2022) 4221: 4231-4249. . **IHC; tested species: mouse**

Striatal Direct Pathway Targets Npas1+ Pallidal Neurons.
Cui Q, Du X, Chang IYM, Pamukcu A, Lilascharoen V, Berceau BL, García D, Hong D, Chon U, Narayanan A, Kim Y, et al.
The Journal of neuroscience : the official journal of the Society for Neuroscience (2021) 4118: 3966-3987. . **IHC; tested species: mouse**

Selected General References

Conditional Wwox deletion in mouse mammary gland by means of two Cre recombinase approaches.
Ferguson BW, Gao X, Kil H, Lee J, Benavides F, Abba MC, Aldaz CM
PloS one (2012) 75: e36618. .

Using the Cre/lox system for targeted integration into the human genome: loxFAS-loxP pairing and delayed introduction of Cre DNA improve gene swapping efficiency.
Lanza AM, Dyess TJ, Alper HS
Biotechnology journal (2012) 77: 898-908. .

Cre recombinase resources for conditional mouse mutagenesis.
Smedley D, Salimova E, Rosenthal N
Methods (San Diego, Calif.) (2011) 534: 411-6. .

Cre/loxP-mediated chromosome engineering of the mouse genome.
Brault V, Besson V, Magnol L, Duchon A, Hérault Y
Handbook of experimental pharmacology (2007) 178: 29-48. .

Bacteriophage P1 site-specific recombination. Purification and properties of the Cre recombinase protein.
Abremski K, Hoess R
The Journal of biological chemistry (1984) 2593: 1509-14. .

The nature of the interaction of the P1 recombinase Cre with the recombining site loxP.
Hoess R, Abremski K, Sternberg N
Cold Spring Harbor symposia on quantitative biology (1984) 49: 761-8. .

Access the online factsheet including applicable protocols
at <https://sysy.com/product/257004> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.