

Aldh1L1

Cat.No. 278 102; Polyclonal rabbit antibody, 200 µl antiserum (lyophilized)

Data Sheet

| | |
|-------------------------|---|
| Reconstitution/ Storage | 200 µl antiserum, lyophilized. For reconstitution add 200 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications | WB: 1 : 1000 up to 1 : 2000 IP: not tested yet ICC: 1 : 500 IHC: 1 : 1000 up to 1 : 2000 IHC_P: 1 : 200 up to 1 : 1000 IHC_G: 1 : 500 up to 1 : 1000 (see remarks) |
| Immunogen | Recombinant protein corresponding to AA 600 to 800 from mouse Aldh1l1 (UniProt Id: Q8R0Y6) |
| Reactivity | Reacts with: rat (P28037), mouse (Q8R0Y6). Other species not tested yet. |
| Specificity | Specific for Aldh1L1 |
| Remarks | IHC_G: Fixation with 3% glyoxal, 1% acetic acid, 20% ethanol in ddH ₂ O according to Richter et al. 2017, or 9% glyoxal, 8% acetic acid in ddH ₂ O according to Konno et al. 2023 are recommended. |

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Aldehyde dehydrogenase family 1 member L1 (**Aldh1L1**), also known as 10-formyltetrahydrofolate dehydrogenase (**FDH**) is a cytosolic enzyme involved in folate metabolism. It has been shown to be involved in the regulation of cell proliferation and is downregulated in malignant human tumors and cancer cell lines.

Aldh1L1 is highly expressed in several cell-types like hepatocytes and astrocytes.

Selected General References

Epigenetic Silencing of ALDH1L1, a Metabolic Regulator of Cellular Proliferation, in Cancers.
Oleinik NV, Krupenko NI, Krupenko SA
Genes & cancer (2011) 22: 130-9. .

Molecular comparison of GLT1+ and ALDH1L1+ astrocytes in vivo in astroglial reporter mice.
Yang Y, Vidensky S, Jin L, Jie C, Lorenzini I, Frankl M, Rothstein JD
Glia (2011) 592: 200-7. .

Gene expression profiling of NF-1-associated and sporadic pilocytic astrocytoma identifies aldehyde dehydrogenase 1 family member L1 (ALDH1L1) as an underexpressed candidate biomarker in aggressive subtypes.
Rodriguez FJ, Giannini C, Asmann YW, Sharma MK, Perry A, Tibbetts KM, Jenkins RB, Scheithauer BW, Anant S, Jenkins S, Eberhart CG, et al.
Journal of neuropathology and experimental neurology (2008) 6712: 1194-204. .

A transcriptome database for astrocytes, neurons, and oligodendrocytes: a new resource for understanding brain development and function.

Cahoy JD, Emery B, Kaushal A, Foo LC, Zamanian JL, Christopherson KS, Xing Y, Lubischer JL, Krieg PA, Krupenko SA, Thompson WJ, et al.

The Journal of neuroscience : the official journal of the Society for Neuroscience (2008) 281: 264-78. .

The folate metabolic enzyme ALDH1L1 is restricted to the midline of the early CNS, suggesting a role in human neural tube defects.

Anthony TE, Heintz N

The Journal of comparative neurology (2007) 5002: 368-83. .

10-formyltetrahydrofolate dehydrogenase, one of the major folate enzymes, is down-regulated in tumor tissues and possesses suppressor effects on cancer cells.

Krupenko SA, Oleinik NV

Cell growth & differentiation : the molecular biology journal of the American Association for Cancer Research (2002) 135: 227-36. .

Access the online factsheet including applicable protocols at <https://susy.com/product/278102> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.