SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

VDAC1

Cat.No. 279 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 1000 (AP staining) **IP**: not tested yet **ICC**: not recommended **IHC**: not recommended **IHC-P (FFPE)**: not tested yet
Immunogen	Synthetic peptide corresponding to AA 15 to 29 from mouse VDAC1 (UniProt Id: Q60932-1)
Reactivity	Reacts with: rat (Q9Z2L0), mouse (Q60932). Other species not tested yet.
Specificity	Due to sequence homologies, probably crossreacts with VDAC 2 and 3
Matching control	279-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Voltage-**d**ependent **a**nion **c**hannels (**VDAC**s), also known as **porins**, are integral membrane proteins which form a channel that allows diffusion of small hydrophilic molecules. Three closely related isoforms, VDAC **1**, 2 and 3 have been described, so far.
VDACs have been originally identified as components of the outer mitochondrial membrane but they also occur in the plasma membrane of different cell types.

Selected General References

Voltage-dependent anion channels (VDACs, porin) expressed in the plasma membrane regulate the differentiation and function of human osteoclasts.
Kotake S et al. Cell Biol. Int. (2013) PubMed:23319323

The expression level of the voltage-dependent anion channel controls life and death of the cell.
Abu-Hamad S et al. Proc. Natl. Acad. Sci. U.S.A. (2006) PubMed:16585511

Bax interacts with the voltage-dependent anion channel and mediates ethanol-induced apoptosis in rat hepatocytes.
Adachi M et al. Am. J. Physiol. Gastrointest. Liver Physiol. (2004) PubMed:15044178

Voltage-dependent anion channel localises to the plasma membrane and peripheral but not perinuclear mitochondria.
Bahamonde MI et al. Pflugers Arch. (2003) PubMed:12698369

Bax interacts with the permeability transition pore to induce permeability transition and cytochrome c release in isolated mitochondria.
Narita M et al. Proc. Natl. Acad. Sci. U.S.A. (1998) PubMed:9843949

Porin pores of mitochondrial outer membranes from high and low eukaryotic cells: biochemical and biophysical characterization.
De Pinto V et al. Biochim. Biophys. Acta (1987) PubMed:3676299

Structure and mode of action of a voltage dependent anion-selective channel (VDAC) located in the outer mitochondrial membrane.
Colombini M et al. Ann. N. Y. Acad. Sci. (1980) PubMed:6249159

Access the online factsheet including applicable protocols at https://sysy.com/product/279003 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.