

## p34-Arc (ARPC2)

Cat.No. 306 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

### Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For <b>reconstitution</b> add 100 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> not recommended <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 <b>IHC:</b> not tested yet <b>IHC-P (FFPE):</b> not tested yet
Clone	334D4
Subtype	IgG1 (κ light chain)
Immunogen	Recombinant protein corresponding to AA 1 to 300 from bovine p34-Arc (UniProt Id: Q3MHR7)
Epitop	AA 229 to 240 from bovine p34-Arc (UniProt Id: Q3MHR7)
Reactivity	Reacts with: human (O15144), cow (Q3MHR7), pig, rat (P85970), mouse (Q9CVB6). Other species not tested yet.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

### Background

Actin polymerization is a necessary event for cell motility. Spontaneous actin oligomerization is slow at given monomeric actin concentrations in cells. The **Arp 2/3 complex** which is about 220 kDa in size has turned out to initiate the polymerization of new actin filaments. This complex consists of two actin like proteins Arp2 and Arp3 and five additional proteins: **p16-Arc** (ArpC5), **p20-Arc** (ArpC4), **p21-Arc** (ArpC3), **p34-Arc** (ArpC2) and p41-Arc (ArpC1). Expression of partial complexes revealed that a heterodimer of p20-Arc and p34-Arc constitutes the core of the complex whereas the remaining subunits are peripherally located.

### Selected References for 306 011

Golgi-localized GAP for Cdc42 functions downstream of ARF1 to control Arp2/3 complex and F-actin dynamics. Dubois T, Paléotti O, Mironov AA, Fraisier V, Stradal TE, De Matteis MA, Franco M, Chavrier P Nature cell biology (2005) 74: 353-64. . **ICC; tested species: human**

Presynapses contain distinct actin nanostructures.

Bingham D, Jakobs CE, Wernert F, Boroni-Rueda F, Jullien N, Schentarra EM, Friedl K, Da Costa Moura J, van Bommel DM, Caillol G, Ogawa Y, et al.

The Journal of cell biology (2023) 22210: . . **ICC; tested species: rat**

### Selected General References

Structural insights into actin-binding, branching and bundling proteins. Winder SJ et al. Curr. Opin. Cell Biol. (2003) PubMed:12517699

Cellular motility driven by assembly and disassembly of actin filaments. Pollard TD et al. Cell (2003) PubMed:12600310

Structure and function of the Arp2/3 complex.

Pollard TD et al. Curr. Opin. Struct. Biol. (2002) PubMed:12504682

Reconstitution of human Arp2/3 complex reveals critical roles of individual subunits in complex structure and activity. Gournier H et al. Mol. Cell (2001) PubMed:11741539

Access the online factsheet including applicable protocols at <https://sysy.com/product/306011> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.