SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

SATB2

Cat.No. 327 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: not recommended **IP**: not tested yet **ICC**: 1 : 500 **IHC**: 1 : 200 **IHC-P (FFPE)**: 1 : 200 up to 1 : 500
Immunogen	Synthetic peptide corresponding to AA 718 to 733 from mouse SATB2 (UniProt Id: Q8VI24)
Reactivity	Reacts with: rat (D3ZJ19), mouse (Q8VI24), monkey. Other species not tested yet.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

The **S**pecial **AT**-rich sequence-**b**inding protein **2** or **SATB2** is a transcription factor required for the initiation of the genetic program for the upper-layer neurons (UL1).
Together with Ctip 2, Coup-TFI, and Fezf2 it is involved in the fine-tuned sequential formation and specification of the different excitatory neuron populations forming the definitive six-layered cortical structure (1-4).
Beyond its pivotal role in neurodevelopment, SATB2 is also expressed in non-neuronal tissues, notably in the epithelium of the lower gastrointestinal tract, including the colon. In normal physiology, it is strongly expressed in glandular cells of the colorectal mucosa. In pathology, SATB2 serves as a highly sensitive and specific immunohistochemical marker for colorectal epithelial differentiation and is expressed in the majority of primary and metastatic colorectal adenocarcinomas (5, 6). This colonic expression pattern makes SATB2 a valuable diagnostic biomarker in surgical pathology, particularly in determining the origin of metastatic lesions.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

Selected References for 327 003

Ageing promotes microglial accumulation of slow-degrading synaptic proteins.
Guldner IH, Wagner VP, Moran-Losada P, Shi SM, Golub SW, Hevler JF, Chen K, Meese BT, Ghoochani A, Pulido E, Oh HS, et al.
Nature (2026) : . . **IHC, IHC-P; tested species: mouse**

Selected General References

Unc5C and DCC act downstream of Ctip2 and Satb2 and contribute to corpus callosum formation.
Srivatsa S et al. Nat Commun (2014) PubMed:24739528

SATB2 as an Immunohistochemical Marker for Colorectal Adenocarcinoma: A Concise Review of Benefits and Pitfalls.
Berg KB et al. Arch Pathol Lab Med (2017) PubMed:28968158

The role of SATB2 as a diagnostic marker for tumors of colorectal origin: Results of a pathology-based clinical prospective study.
Dragomir A et al. Am J Clin Pathol (2014) PubMed:24713733

The CB(1) cannabinoid receptor drives corticospinal motor neuron differentiation through the Ctip2/Satb2 transcriptional regulation axis.
Díaz-Alonso J et al. J. Neurosci. (2012) PubMed:23175820

The sympathetic neurotransmitter switch depends on the nuclear matrix protein Satb2.
Apostolova G et al. J. Neurosci. (2010) PubMed:21123581

SATB2 interacts with chromatin-remodeling molecules in differentiating cortical neurons.
Gyorgy AB et al. Eur. J. Neurosci. (2008) PubMed:18333962

Access the online factsheet including applicable protocols at https://sysy.com/product/327003 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.