

DOPA decarboxylase

Cat.No. 369 011; Monoclonal mouse antibody, 100 µg purified IgG (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µg purified IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 100 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 IP: not tested yet ICC: not tested yet IHC: 1 : 500 up to 1 : 5000 IHC-P/FFPE: 1 : 1000
Clone	126F10
Subtype	IgG2b (κ light chain)
Immunogen	Recombinant protein corresponding to AA 1 to 480 from mouse DDC (UniProt Id: O88533)
Reactivity	Reacts with: mouse (O88533), rat (P14173). Other species not tested yet.
Specificity	Specific for DOPA decarboxylase

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Access the online factsheet including applicable protocols at <https://sysy.com/product/369011> or scan the QR-code.



Background

DOPA decarboxylase, also referred to as **aromatic L-amino acid decarboxylase, AADC** or **DDC**, is the second enzyme in the biosynthesis of dopamine and serotonin. In addition, it is responsible for the synthesis of the trace amines phenylethylamine, p-tyramine, and tryptamine, which are considered to act as neuromodulators.

DDC is expressed in the central nervous system, but has also been detected in several tissues such as liver, kidney, small intestine, adrenal gland and blood vessels.

The inhibition of DDC is used for the treatment of Parkinson's disease.

Selected General References

Marked disparity between age-related changes in dopamine and other presynaptic dopaminergic markers in human striatum.
Haycock JW, Becker L, Ang L, Furukawa Y, Hornykiewicz O, Kish SJ
Journal of neurochemistry (2003) 873: 574-85. .

Structural insight into Parkinson's disease treatment from drug-inhibited DOPA decarboxylase.
Burkhard P, Dominici P, Borri-Voltattorni C, Jansonius JN, Malashkevich VN
Nature structural biology (2001) 811: 963-7. .

Aromatic L-amino acid decarboxylase: a neglected and misunderstood enzyme.
Berry MD, Juorio AV, Li XM, Boulton AA
Neurochemical research (1996) 219: 1075-87. .

Functionally important residues of aromatic L-amino acid decarboxylase probed by sequence alignment and site-directed mutagenesis.

Ishii S, Mizuguchi H, Nishino J, Hayashi H, Kagamiyama H
Journal of biochemistry (1996) 1202: 369-76. .

Distinct promoters direct neuronal and nonneuronal expression of rat aromatic L-amino acid decarboxylase.
Albert VR, Lee MR, Bolden AH, Wurzburger RJ, Aguanno A
Proceedings of the National Academy of Sciences of the United States of America (1992) 8924: 12053-7. .

Aromatic L-amino acid decarboxylase in the rat brain: immunocytochemical localization in neurons of the brain stem.
Jaeger CB, Ruggiero DA, Albert VR, Park DH, Joh TH, Reis DJ
Neuroscience (1984) 113: 691-713. .

FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.