

## Numblike

Cat.No. 374 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 100 <b>IHC:</b> not tested yet <b>IHC-P:</b> not tested yet <b>IHC-Fr:</b> 1 : 500 (see remarks)
Immunogen	Synthetic peptide corresponding to AA 579 to 595 from mouse Numbl (UniProt Id: O08919)
Reactivity	Reacts with: rat (A1L1I3), mouse (O08919). Other species not tested yet.
Specificity	K.O. validated
Matching control	374-0P
Remarks	<b>IHC-Fr:</b> 4% formaldehyde/PFA fixation is recommended.

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

Numb proteins (**Numbl**ike and Numb) display a complex pattern of functions such as the control of asymmetric cell division, cell fate choice, endocytosis, cell adhesion, and cell migration. They have been shown to inhibit Notch signaling by stimulating endocytosis of Notch. Numb and Numbl like have at least partially distinct functions. Numbl like is a negative regulator of the NF-κB signaling pathway by abrogating TRAF5-induced activation of NF-κB. Recently, Numbl like was implicated as a physiologically relevant target of microRNA miR-34a in neural progenitor cells allowing for enhanced Notch signaling and inhibition of neuronal differentiation. Numb and Numbl like are essential in maintaining neural progenitor cells during early neurogenesis by allowing cells to choose progenitor over neuronal fates. They were recently also discovered to be involved in cardiac morphogenesis.

## Selected General References

- Numb family proteins: novel players in cardiac morphogenesis and cardiac progenitor cell differentiation. Wu M et al. *Biomol Concepts* (2015) PubMed:25883210
- Precardiac deletion of Numb and Numbl like reveals renewal of cardiac progenitors. Shenje LT et al. *Elife* (2014) PubMed:24843018
- Numbl like regulates proliferation, apoptosis, and invasion of lung cancer cell. Yingjie L et al. *Tumour Biol.* (2013) PubMed:23681800
- Inhibition of Notch2 by Numb/Numbl like controls myocardial compaction in the heart. Yang J et al. *Cardiovasc. Res.* (2012) PubMed:22865640
- MiR-34a represses Numbl in murine neural progenitor cells and antagonizes neuronal differentiation. Fineberg SK et al. *PLoS ONE* (2012) PubMed:22701667
- Numbl inhibits glioma cell migration and invasion by suppressing TRAF5-mediated NF-κB activation. Tao T et al. *Mol. Biol. Cell* (2012) PubMed:22593207
- Numbl like and Numb differentially affect p53 and Sonic Hedgehog signaling. Liu L et al. *Biochem. Biophys. Res. Commun.* (2011) PubMed:21893032
- The multiple functions of Numb. Gulino A et al. *Exp. Cell Res.* (2010) PubMed:19944684
- Numb and Numbl are required for maintenance of cadherin-based adhesion and polarity of neural progenitors. Rasin MR et al. *Nat. Neurosci.* (2007) PubMed:17589506
- Postnatal deletion of Numb/Numbl like reveals repair and remodeling capacity in the subventricular neurogenic niche. Kuo CT et al. *Cell* (2006) PubMed:17174898
- Continuing role for mouse Numb and Numbl in maintaining progenitor cells during cortical neurogenesis. Petersen PH et al. *Nat. Neurosci.* (2004) PubMed:15273690
- Differential expression of mammalian Numb, Numbl like and Notch1 suggests distinct roles during mouse cortical neurogenesis. Zhong W et al. *Development* (1997) PubMed:9169836

Access the online factsheet including applicable protocols at <https://sysy.com/product/374003> or scan the QR-code.



# FAQ - How should I store my antibody?

## Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

## Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

## Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

## Product Specific Hints for Storage

### Control proteins / peptides

- Store at -20°C to -80°C

### Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

### Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

# Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.