

HSP90 β

Cat.No. 380-1P; control peptide, 100 μ g peptide (lyophilized)

Data Sheet

Reconstitution/ Storage	100 μ g peptide, lyophilized. For reconstitution add 100 μ l H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Control peptides should be stored at -20°C when still lyophilized! For detailed information, see back of the data sheet.
Immunogen	Synthetic peptide corresponding to AA 639 to 709 from human HSP90 β (UniProt Id: P08238)
Recommended dilution	Optimal concentrations should be determined by the end-user.
Matching antibodies	380 103
Remarks	This control peptide consists of the synthetic peptide (aa 693-709 of human HSP 90 β) that has been used for immunization. It has been tested in preadsorption experiments and blocks efficiently and specifically the corresponding signal in Western blots. The amount of peptide needed for efficient blocking depends on the titer and on the affinity of the antibody to the antigen.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Heat shock proteins 90 (HSP 90) are molecular chaperones. Two isoforms (α and β) have been described so far. They promote the maturation, structural maintenance and proper regulation of specific target proteins involved for instance in cell cycle control and signal transduction. They undergo a functional cycle that is linked to its ATPase activity. This cycle probably induces conformational changes in the client proteins, thereby causing their activation. HSP 90s interact dynamically with various co-chaperones that modulate their substrate recognition, ATPase cycle and chaperone function.

Selected General References

From Conformation to Interaction: Techniques to Explore the Hsp70/Hsp90 Network.

Batista FA, Gava LM, Pinheiro GM, Ramos CH, Borges JC
Current protein & peptide science (2015) 168: 735-53. .

Hsp90, the concertmaster: tuning transcription.

Khurana N, Bhattacharyya S
Frontiers in oncology (2015) 5: 100. .

The cellular world according to Hsp90.

Richter K, Hendershot LM, Freeman BC
Nature structural & molecular biology (2007) 142: 90-4. .

Analysis of native forms and isoform compositions of the mouse 90-kDa heat shock protein, HSP90.

Minami Y, Kawasaki H, Miyata Y, Suzuki K, Yahara I
The Journal of biological chemistry (1991) 266: 10099-103. .

Stage and lineage-regulated expression of two hsp90 transcripts during mouse germ cell differentiation and embryogenesis.

Gruppi CM, Zakeri ZF, Wolgemuth DJ
Molecular reproduction and development (1991) 283: 209-17. .

Mapping the HSP90 binding region of the glucocorticoid receptor.

Howard KJ, Holley SJ, Yamamoto KR, Distelhorst CW
The Journal of biological chemistry (1990) 265: 11928-35. .

Two mammalian heat shock proteins, HSP90 and HSP100, are actin-binding proteins.

Koyasu S, Nishida E, Kadowaki T, Matsuzaki F, Iida K, Harada F, Kasuga M, Sakai H, Yahara I
Proceedings of the National Academy of Sciences of the United States of America (1986) 83: 8054-8. .

Access the online factsheet including applicable protocols at <https://sysy.com/product/380-1P> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.