

## Spinophilin

Cat.No. 399 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

### Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin was added for stabilization. For <b>reconstitution</b> add 50 µl H <sub>2</sub> O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	<b>WB:</b> 1 : 1000 (AP staining) <b>IP:</b> not tested yet <b>ICC:</b> 1 : 500 up to 1 : 1000 <b>IHC:</b> 1 : 500 up to 1 : 1000 <b>IHC-P:</b> 1 : 500 <b>DNA-PAINT:</b> yes (see remarks)
Immunogen	Synthetic peptide corresponding to AA 290 to 302 from rat Spinophilin (UniProt Id: O35274)
Reactivity	Reacts with: rat (O35274), mouse (Q6R891). Other species not tested yet.
Specificity	Specific for Spinophilin isoform1. Does not detect isoform 2. K.O. validated
Matching control	399-0P
Remarks	<b>DNA-PAINT:</b> This antibody has been successfully used for DNA-PAINT application (see Unterauer et al., 2024; <a href="https://pubmed.ncbi.nlm.nih.gov/38552614/">PMID: 38552614</a> ).

**TO BE USED IN VITRO / FOR RESEARCH ONLY**  
**NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS**

## Background

**Spinophilin**, also known as **Neurabin II**, is a scaffolding protein enriched in neuronal dendritic spines. It is involved in the targeted localization of protein phosphatase-1 (PP1), critical for dopaminergic modulation of glutamate neurotransmission. More recently it has also been identified as a tumor suppressor. Loss of its function induces a proliferative response by increasing retinoblastoma protein (pRb) phosphorylation.

## Selected References for 399 003

Spatial proteomics in neurons at single-protein resolution.  
Unterauer EM, Shetab Boushehri S, Jevdokimenko K, Masullo LA, Ganji M, Sograte-Idrissi S, Kowalewski R, Strauss S, Reinhardt SCM, Perovic A, Marr C, et al.  
Cell (2024) 1877: 1785-1800.e16. . **DNA\_PAINT; tested species: rat**

Protocol for SUM-PAINT spatial proteomic imaging generating neuronal architecture maps in rat hippocampal neurons.  
Unterauer EM, Schentarra EM, Jevdokimenko K, Boushehri SS, Marr C, Opazo F, Fornasiero EF, Jungmann R  
STAR protocols (2025) 61: 103637. . **DNA\_PAINT; tested species: rat**

## Selected General References

Characterisation of spinophilin immunoreactivity in postmortem human brain homogenates.  
Brocos-Mosquera I et al. Prog. Neuropsychopharmacol. Biol. Psychiatry (2018) PubMed:28941770

Spinophilin: a new tumor suppressor at 17q21.  
Carnero A et al. Curr. Mol. Med. (2012) PubMed:22515982

The identification of a second actin-binding region in spinophilin/neurabin II.  
Barnes AP et al. Brain Res. Mol. Brain Res. (2004) PubMed:15135218

Phosphorylation of spinophilin modulates its interaction with actin filaments.  
Hsieh-Wilson LC et al. J. Biol. Chem. (2003) PubMed:12417592

Characterization of the neuronal targeting protein spinophilin and its interactions with protein phosphatase-1.  
Hsieh-Wilson LC et al. Biochemistry (1999) PubMed:10194355

Neurabin-II/spinophilin. An actin filament-binding protein with one pdz domain localized at cadherin-based cell-cell adhesion sites.  
Satoh A et al. J. Biol. Chem. (1998) PubMed:9452470

Spinophilin, a novel protein phosphatase 1 binding protein localized to dendritic spines.  
Allen PB et al. Proc. Natl. Acad. Sci. U.S.A. (1997) PubMed:9275233

Access the online factsheet including applicable protocols at <https://sysy.com/product/399003> or scan the QR-code.



## FAQ - How should I store my antibody?

### Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

### Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

### Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

### Product Specific Hints for Storage

#### Control proteins / peptides

- Store at -20°C to -80°C.

#### Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

#### Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

## Avoid repeated freeze-thaw cycles for all antibodies!

## FAQ - How should I reconstitute my antibody?

### Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.