SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

TMEM119

Cat.No. 400 203; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: not tested yet **IP**: not tested yet **ICC**: not tested yet **IHC**: 1 : 500 up to 1 : 1000 **IHC-P (FFPE)**: not recommended
Immunogen	Recombinant protein corresponding to AA 189 to 283 from rat TMEM119 (UniProt Id: B2RYL3)
Reactivity	Reacts with: rat (B2RYL3). Weaker signal: mouse (Q8R138). Other species not tested yet.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Microglia are resident myeloid cells of the central nervous system (CNS). They are ontogenetically and functionally distinct from monocyte-derived macrophages that infiltrate the CNS under pathological conditions. **T**rans**mem**brane protein **119** (**TMEM119**) is a single-pass type I membrane protein that has been identified as a useful, highly selective microglia marker protein.

Selected References for 400 203

Immunohistological and Ultrastructural Study of the Inflammatory Response to Perforated Polyimide Cortical Implants: Mechanisms Underlying Deterioration of Electrophysiological Recording Quality.
Huang SH, Shmoel N, Jankowski MM, Erez H, Sharon A, Abu-Salah W, Nelken I, Weiss A, Spira ME
Frontiers in neuroscience (2020) 14: 926. . **IHC; tested species: rat**

Microglia regulate neuronal activity via structural remodeling of astrocytes.
Gu N, Makashova O, Laporte C, Chen CQ, Li B, Chevillard PM, Lean G, Yang J, Wong C, Fan J, Sharif B, et al.
Neuron (2025) 11320: 3408-3423.e5. . **IHC; tested species: rat**

Significant Sex Differences in the Efficacy of the CSF1R Inhibitor-PLX5622 on Rat Brain Microglia Elimination.
Sharon A, Erez H, Spira ME
Pharmaceuticals (Basel, Switzerland) (2022) 155: . . **IHC; tested species: rat**

Inflammatory Foreign Body Response Induced by Neuro-Implants in Rat Cortices Depleted of Resident Microglia by a CSF1R Inhibitor and Its Implications.
Sharon A, Jankowski MM, Shmoel N, Erez H, Spira ME
Frontiers in neuroscience (2021) 15: 646914. . **IHC; tested species: rat**

Selected General References

New tools for studying microglia in the mouse and human CNS.
Bennett ML et al. Proc. Natl. Acad. Sci. U.S.A. (2016) PubMed:26884166

TMEM119 marks a subset of microglia in the human brain.
Satoh J et al. Neuropathology (2016) PubMed:26250788

Access the online factsheet including applicable protocols at https://sysy.com/product/400203 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.