

Oxytocin

Cat.No. 408 008; Recombinant rabbit antibody, 50 µg recombinant IgG (lyophilized)

Data Sheet

| | |
|----------------------------|---|
| Reconstitution/ Storage | 50 µg purified recombinant IgG, lyophilized. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet. |
| Applications | WB: not tested yet Dot blot: 1 : 5000 up to 1 : 20000 (AP staining) (see remarks) IP: not tested yet ICC: not tested yet IHC: 1 : 1000 up to 1 : 5000 IHC-P (FFPE): 1 : 1000 |
| Clone | Rb323B2 |
| Subtype | IgG1 (κ light chain) |
| Immunogen | Synthetic peptide corresponding to the processed bioactive oxytocin peptide cleaved from mouse oxytocin-neurophysin1 (UniProt Id: P35454) |
| Reactivity | Reacts with: mouse (P35454), rat (P01179). Other species not tested yet. |
| Specificity | The antibody preferentially recognizes the processed bioactive oxytocin peptide. The unprocessed precursor protein is only recognized to a limited extent, if at all. The antibody does not crossreact with vasopressin. |
| Remarks | This antibody is a chimeric antibody based on the monoclonal rat antibody SY-323B2. The constant regions of the heavy and light chains have been replaced with rabbit specific sequences. Therefore, the antibody can be used with standard anti-rabbit secondary reagents. The antibody has been expressed in mammalian cells. Dot blot: This application was only tested with synthetic peptides. |

Background

Oxytocin is a peptide hormone which is synthesized as an inactive precursor in nerve cell bodies in the supraoptic nucleus and paraventricular nucleus of the hypothalamus. The precursor protein is progressively hydrolyzed to produce oxytocin and its carrier protein neurophysin I. Together with neurophysin I, it is packaged into neurosecretory vesicles and transported axonally to the nerve endings in the neurohypophysis, where it is either stored or secreted into the bloodstream. Oxytocin is also produced by some neurons of the hypothalamus that project to other parts of the brain and to the spinal cord. Oxytocin is best known for roles in female reproduction as its release causes smooth muscle contraction during parturition and lactation. It is also involved in cognition, tolerance, adaptation, and complex sexual and maternal behavior. Oxytocin is used as a medication to facilitate childbirth.

Selected General References

Oxytocin and vasopressin: linking pituitary neuropeptides and their receptors to social neurocircuits. Baribeau DA et al. *Front Neurosci* (2015) PubMed:26441508

Neuromodulation by oxytocin and vasopressin in the central nervous system as a basis for their rapid behavioral effects. Stoop R et al. *Curr. Opin. Neurobiol.* (2014) PubMed:25463629

Oxytocin modulates female sociosexual behavior through a specific class of prefrontal cortical interneurons. Nakajima M et al. *Cell* (2014) PubMed:25303526

Oxytocin: the great facilitator of life. Lee HJ et al. *Prog. Neurobiol.* (2009) PubMed:19482229

Access the online factsheet including applicable protocols at <https://sysy.com/product/408008> or scan the QR-code.



TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.