

Glutaminase 1

Cat.No. 456 003; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For reconstitution add 50 µl H ₂ O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP-staining) IP: not tested yet ICC: 1 : 500 up to 1 : 1000 IHC: 1 : 1000 IHC-P: not tested yet
Immunogen	Synthetic peptide corresponding to AA 655 to 674 from rat Glutaminase (UniProt Id: P13264)
Reactivity	Reacts with: mouse (D3Z7P3), rat (P13264). Other species not tested yet.
Matching control	456-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Glutamate is the major excitatory neurotransmitter in the central nervous system. During neurotransmission it is released from synaptic vesicles into the synaptic cleft and is sensed by post-synaptic glutamate receptors (GluAs) (1).

Excitatory **a**mino **a**cid **t**ransporters (EAATs) translocate extracellular glutamate into neighbouring astrocytes where it is converted to glutamine by glutamine synthetase (2, 3, 4).

The glutamine is shuttled back to neurons where it is processed to glutamate by glutaminases (GLS) (5). This postulated Glu/Gln cycle is a key mechanism for the homeostatic control of these amino acids. In humans, three isoforms (GLS1, 2 and 3) have been described. GLS1 is highly expressed in brain and kidney whereas GLS2 is mainly expressed in cardiac and skeletal muscle. GLS3 shows a broader tissue distribution (5). The GLS1 precursor is processed to a 65 and 68 kDa chain.

Selected General References

Overview of glutamatergic neurotransmission in the nervous system.
Niciu MJ et al. Pharmacol Biochem Behav (2012) PubMed:21889952

Astrocyte glutamine synthetase: pivotal in health and disease.
Rose CF et al. Biochem. Soc. Trans. (2013) PubMed:24256247

The density of EAAC1 (EAAT3) glutamate transporters expressed by neurons in the mammalian CNS.
Holmseth S et al. J. Neurosci. (2012) PubMed:22539860

Cloning and analysis of unique human glutaminase isoforms generated by tissue-specific alternative splicing.
Elgadi KM et al. Physiol Genomics (1999) PubMed:11015561

Synaptic activation of glutamate transporters in hippocampal astrocytes.
Bergles DE et al. Neuron (1997) PubMed:9427252

Access the online factsheet including applicable protocols at <https://sysis.com/product/456003> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All our antibodies and control proteins / peptides are shipped lyophilized (vacuum freeze-dried) and are stable in this form without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at 4°C before reconstitution. **They must not be stored in the freezer when still lyophilized!** Temperatures below zero may cause loss of performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long term storage (several months) may lead to aggregation.
- **Control peptides** should be kept at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- The storage freezer must not be of the frost-free variety ("no-frost freezer"). This cycle between freezing and thawing (to reduce frost-build-up), which is exactly what should be avoided. For the same reason, antibody vials should be placed in an area of the freezer that has minimal temperature fluctuations, for instance towards the back rather than on a door shelf.
- Aliquot the antibody and store frozen (-20°C to -80°C). Avoid very small aliquots (below 20 µl) and use the smallest storage vial or tube possible. The smaller the aliquot, the more the stock concentration is affected by evaporation and adsorption of the antibody to the surface of the storage vial or tube. Adsorption of the antibody to the surface leads to a substantial loss of activity.
- The addition of glycerol to a final concentration of 50% lowers the freezing point of your stock and keeps your antibody at -20°C in liquid state. This efficiently avoids freeze and thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C.

Monoclonal Antibodies

- **Ascites** and **hybridoma supernatant** should be stored at -20°C up to -80°C. **Prolonged storage at 4°C is not recommended!** Unlike serum, ascites may contain proteases that will degrade the antibodies.
- **Purified IgG** should be stored at -20°C up to -80°C. Adding a carrier protein like BSA will increase long term stability. Many of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Polyclonal Antibodies

- **Crude antisera:** With anti-microbials added, they may be stored at 4°C. However, frozen storage (-20°C up to -80°C) is preferable.
- **Affinity purified antibodies:** Less robust than antisera. Storage at -20°C up to -80°C is recommended. Adding a carrier protein like BSA will increase long term stability. Most of our antibodies already contain carrier proteins. Please refer to the data-sheet for detailed information.

Fluorescence-labeled Antibodies

- Store as a liquid with 1 : 1 (v/v) glycerol at -20°C. Protect these antibodies from light exposure.

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All our purified antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the amount of deionized water given in the respective datasheet. If higher volumes are preferred, add water as mentioned above and then the desired amount of PBS and a stabilizing carrier protein (e.g. BSA) to a final concentration of 2%. Some of our antibodies already contain albumin. Take this into account when adding more carrier protein. For complete reconstitution, carefully remove the lid. After adding water, briefly vortex the solution. You can spin down the liquid by placing the vial into a 50 ml centrifugation tube filled with paper.
- If desired, add small amounts of azide or thimerosal to prevent microbial growth. This is especially recommended if you want to keep an aliquot a 4°C.
- After reconstitution of fluorescence-labeled antibodies, add 1 : 1 (v/v) glycerol to a final concentration of 50%. This lowers the freezing point of your stock and keeps your antibody in liquid state at -20°C.
- Glycerol may also be added to unlabeled primary antibodies. It is a suitable way to avoid freeze-thaw cycles.
- Please refer to our **tips and hints for subsequent storage** of reconstituted antibodies and control peptides and proteins.