

Glutaminase1

Cat.No. 456 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 (AP staining) IP: not tested yet ICC: 1 : 1000 up to 1 : 2000 IHC: 1 : 500 up to 1 : 1000 IHC-P: not tested yet
Immunogen	Synthetic peptide corresponding to AA 655 to 674 from rat Glutaminase (UniProt Id: P13264)
Reactivity	Reacts with: mouse (D3Z7P3), rat (P13264). Other species not tested yet.
Matching control	456-0P

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Glutamate is the major excitatory neurotransmitter in the central nervous system. During neurotransmission it is released from synaptic vesicles into the synaptic cleft and is sensed by post-synaptic glutamate receptors (GluRs) (1).

Excitatory **a**mino **a**cid **t**ransporters (EAATs) translocate extracellular glutamate into neighbouring astrocytes where it is converted to glutamine by glutamine synthetase (2, 3, 4).

The glutamine is shuttled back to neurons where it is processed to glutamate by glutaminases (GLS) (5). This postulated Glu/Gln cycle is a key mechanism for the homeostatic control of these amino acids. In humans, three isoforms (GLS1, 2 and 3) have been described. GLS1 is highly expressed in brain and kidney whereas GLS2 is mainly expressed in cardiac and skeletal muscle. GLS3 shows a broader tissue distribution (5). The GLS1 precursor is processed to a 65 and 68 kDa chain.

For more information on protein expression pattern, please refer to the overview image in our SYSY Antibodies ATLAS.

Selected General References

Overview of glutamatergic neurotransmission in the nervous system.
Niciu MJ et al. Pharmacol Biochem Behav (2012) PubMed:21889952

Astrocyte glutamine synthetase: pivotal in health and disease.
Rose CF et al. Biochem. Soc. Trans. (2013) PubMed:24256247

The density of EAAC1 (EAAT3) glutamate transporters expressed by neurons in the mammalian CNS.
Holmseth S et al. J. Neurosci. (2012) PubMed:22539860

Cloning and analysis of unique human glutaminase isoforms generated by tissue-specific alternative splicing.
Elgadi KM et al. Physiol Genomics (1999) PubMed:11015561

Synaptic activation of glutamate transporters in hippocampal astrocytes.
Bergles DE et al. Neuron (1997) PubMed:9427252

Access the online factsheet including applicable protocols at <https://susy.com/product/456004> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
Adding a carrier protein (e.g., BSA) enhances long-term stability.
Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
Adding a carrier protein such as BSA improves long-term stability.
Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.