

JAM-A

Cat.No. 522 004; Polyclonal Guinea pig antibody, 100 µl antiserum (lyophilized)

Data Sheet

Reconstitution/ Storage	100 µl antiserum, lyophilized. For reconstitution add 100 µl H ₂ O, then aliquot and store at -20°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	WB: 1 : 1000 up to 1 : 2000 (AP staining) IP: not tested yet ICC: 1 : 500 (see remarks) IHC: not tested yet IHC-P (FFPE): not tested yet
Immunogen	Recombinant protein corresponding to extracellular-domain of rat JAM-A (UniProt Id: Q9JHY1)
Reactivity	Reacts with: rat (Q9JHY1), mouse (O88792), dog. Other species not tested yet.
Remarks	ICC: The following fixatives are possible: 3% glyoxal, 4% formaldehyde/PFA.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Junctional adhesion molecule-A (JAM-A) is a key transmembrane protein in cell adhesion, immune response, and barrier integrity. It localizes at tight junctions, interacting with ZO-1 and PAR-3 to regulate cell polarity and maintain the blood-brain barrier (BBB) (1). JAM-A also facilitates leukocyte adhesion and transmigration, crucial for immune defense but implicated in inflammatory diseases (1). In the BBB, its downregulation correlates with barrier breakdown in conditions like multiple sclerosis (2). Overexpression in glioblastomas links JAM-A to tumor stem cell maintenance and therapy resistance (3). As a therapeutic target, JAM-A inhibition can reduce leukocyte infiltration and tissue damage, with implications for inflammatory bowel disease, stroke, and cancer (1). Its dual role in junction stabilization and disease progression makes JAM-A a key focus for medical research. JAM-A has a molecular weight of 29 kD which can reach between 32 kD and 43 kD as a result of post-translational modifications (N-glycosylation) or due to differences in mRNA splicing (4).

Selected General References

Tight junction dynamics: the role of junctional adhesion molecules (JAMs).
Garrido-Urbani S et al. Cell Tissue Res (2014) PubMed:24595739

F11R/JAM-A: why do platelets express a molecule which is also present in tight junctions?
Kamola P et al. Platelets (2023) PubMed:37246517

Expression and prognostic value of JAM-A in gliomas.
Rosager AM et al. J Neurooncol (2017) PubMed:28677106

Decreased junctional adhesion molecule-A expression during blood-brain barrier breakdown.
Yeung D et al. Acta Neuropathol (2008) PubMed:18357461

Access the online factsheet including applicable protocols at <https://sysy.com/product/522004> or scan the QR-code.



FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies.** Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free (“no-frost”) freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at -20°C to -80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C. Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C. Adding a carrier protein (e.g., BSA) enhances long-term stability. Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C. Adding a carrier protein such as BSA improves long-term stability. Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at -20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.