SYSY Antibodies

Rudolf-Wissell-Str. 28a
37079 Göttingen, Germany
Phone: +49 551-50556-0
Fax: +49 551-50556-384
E-mail: sales@sysy.com
Web: www.sysy.com

Siglec-10

Cat.No. 556 013; Polyclonal rabbit antibody, 50 µg specific antibody (lyophilized)

Data Sheet

Reconstitution/ Storage	50 µg specific antibody, lyophilized. Affinity purified with the immunogen. Albumin and azide were added for stabilization. For **reconstitution** add 50 µl H_2O to get a 1mg/ml solution in PBS. Then aliquot and store at -20°C to -80°C until use. Antibodies should be stored at +4°C when still lyophilized. Do not freeze! For detailed information, see back of the data sheet.
Applications	**WB**: 1 : 1000 (fluorescent detection) **IP**: not tested yet **ICC**: not tested yet **IHC**: not tested yet **IHC-P (FFPE)**: 1 : 1000 (see remarks)
Immunogen	Recombinant protein corresponding to residues near the carboxy region of human Siglec-10 (UniProt Id: Q96LC7)
Reactivity	Reacts with: human (Q96LC7). Other species not tested yet.
Remarks	**IHC-P (FFPE)**: For best results in IHC-P staining, we recommend 566 003.

TO BE USED IN VITRO / FOR RESEARCH ONLY
NOT TOXIC, NOT HAZARDOUS, NOT INFECTIOUS, NOT CONTAGIOUS

Background

Siglec-10 (sialic acid-binding Ig-like lectin 10) is a member of the CD33-related Siglec family, a subgroup of type I transmembrane receptors characterized by extracellular immunoglobulin-like domains and cytoplasmic immunoreceptor-based tyrosine-based inhibitory motifs (ITIMs), which recruit the phosphatases SHP-1 and SHP-2 to attenuate activating immune signals (1). In humans, Siglec-10 is expressed in a variety of immune cell populations, including tumor-associated macrophages (TAMs), dendritic cells, natural killer (NK) cells, and subsets of activated T and B lymphocytes, with expression levels reportedly increasing significantly within the tumor microenvironment (2–4). The binding of Siglec-10 to its ligand CD24 triggers inhibitory signaling pathways that suppress the activation of T-cell receptors and NK cells, reduce cytokine production, and impair macrophage-mediated phagocytosis, thereby contributing to the establishment of a “don’t-eat-me” immune checkpoint (2).
Clinically, elevated Siglec-10 expression has been associated with poorer overall survival and independent prognostic significance, likely due to the accumulation of Siglec-10 tumor-associated macrophages (TAMs) and their role in reshaping the tumor immune microenvironment (3-5).

Selected General References

Unravelling CD24-Siglec-10 pathway: Cancer immunotherapy from basic science to clinical studies.
Hazra R et al. Immunology (2024) PubMed:39129256

High SIGLEC10-Expressing Tumor-Associated Macrophages Participate in Remodeling the Tumor Immune Microenvironment and Predict Poor Prognosis in Colorectal Cancer.
Miao C et al. Cancer Med (2026) PubMed:42487207

Targeting Interactions Between Siglec-10 and α3β1 Integrin Enhances Macrophage-Mediated Phagocytosis of Pancreatic Cancer.
Saini P et al. Cancer Res (2026) PubMed:41182080

High SIGLEC10-Expressing Tumor-Associated Macrophages Participate in Remodeling the Tumor Immune Microenvironment and Predict Poor Prognosis in Colorectal Cancer.
Miao C et al. Cancer Med (2026) PubMed:42487207

Molecular Mechanism of Tumor Cell Immune Escape Mediated by CD24/Siglec-10.
Yin SS et al. Front Immunol (2020) PubMed:32765491

Access the online factsheet including applicable protocols at https://sysy.com/product/556013 or scan the QR-code.

FAQ - How should I store my antibody?

Shipping Conditions

- All SYSY antibodies and control proteins/peptides are shipped lyophilized (vacuum freeze-dried). In this form, they remain stable without loss of quality at ambient temperatures for several weeks.

Storage of Sealed Vials after Delivery

- **Unlabeled** and **biotin-labeled antibodies** and **control proteins** should be stored at **4°C** before reconstitution. **Do not freeze lyophilized antibodies**. Temperatures below 0°C may impair performance.
- **Fluorescence-labeled antibodies** should be reconstituted immediately upon receipt. Long-term storage of lyophilized fluorophore-conjugates may cause aggregation.
- **Control peptides** should be stored at -20°C before reconstitution.

Long Term Storage after Reconstitution (General Considerations)

- **Do not use frost-free ("no-frost") freezers.** These units periodically warm to remove ice buildup, causing freeze–thaw cycles that can damage antibodies.
- Store vials in areas with minimal temperature fluctuation - preferably toward the back of the freezer, not on the door.
- Aliquot reconstituted antibodies and store at –20°C to –80°C.
- Avoid very small aliquots (<20 µL), as evaporation and adsorption to tube surfaces can reduce antibody concentration and activity.
- Use the smallest practical storage vial to minimize surface area.
- Adding glycerol to a final concentration of 50% prevents freezing at -20°C, allowing storage in liquid form and effectively avoiding freeze–thaw cycles.

Product Specific Hints for Storage

Control proteins / peptides

- Store at -20°C to -80°C

Monoclonal Antibodies

- **Ascites and hybridoma supernatant:** Store at -20°C to -80°C
  Prolonged storage at 4°C is not recommended, as proteases present in ascites may degrade antibodies.
- **Purified IgG:** Store at -20°C to -80°C
  Adding a carrier protein (e.g., BSA) enhances long-term stability.
  Many SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Polyclonal Antibodies

- **Crude antisera:** Can be stored at 4°C with antimicrobials added, but -20°C to -80°C is preferred
- **Affinity-purified antibodies:** Less stable than antisera; store at -20°C to -80°C
  Adding a carrier protein such as BSA improves long-term stability.
  Most SYSY antibodies already contain carrier proteins - refer to the respective datasheet for details.

Fluorescence-labeled Antibodies

- Store as a liquid with 1:1 (v/v) glycerol at -20°C, and protect from light exposure

Avoid repeated freeze-thaw cycles for all antibodies!

FAQ - How should I reconstitute my antibody?

Reconstitution

- All purified SYSY antibodies are lyophilized from PBS. To reconstitute the antibody in PBS, add the volume of deionized water specified in the corresponding datasheet. If a larger final volume is desired, first add the recommended amount of water, then adjust with PBS and, if needed, add a stabilizing carrier protein (e.g., BSA) to a final concentration of 2%. Some SYSY antibodies already contain albumin; please take this into account before adding additional carrier protein.

For complete reconstitution, carefully remove the vial cap. After adding water, briefly vortex the solution. To collect the liquid at the bottom of the vial, place the vial inside a 50 ml centrifuge tube padded with paper and centrifuge briefly.

- If desired, small amounts of azide or thimerosal may be added to prevent microbial growth. This is particularly recommended when storing an aliquot at 4°C.
- After reconstitution of fluorescence-labeled antibodies, add glycerol 1:1 (v/v) to achieve a final concentration of 50%. This prevents freezing at –20°C and keeps the antibody in liquid form, effectively avoiding freeze–thaw cycles.
- Glycerol may also be added to unlabeled primary antibodies as a general measure to prevent freeze–thaw damage.
- For further guidance, please refer to our **storage tips** and recommendations for reconstituted antibodies, control peptides, and control proteins.