

## Protocol for mCLING (Cat. No. 710 006AT488)

### Immunohistochemistry (IHC)

### Membrane Staining - mCLING

#### General considerations

mCLING can also be used to study synaptic vesicle recycling in more complex tissue preparations. Of course, tissue dissection and treatment must maintain the sample in a viable state to ensure the ongoing of membrane recycling events.

The main practical difficulty when labeling tissues is ensuring that mCLING homogeneously reaches the plasma membrane of the cell or structure of interest. Therefore, it is recommended to make preliminary tests with different incubation times and concentrations. Once efficient penetration is ensured, preparations can be electrically stimulated or depolarized with a solution containing a high concentration of  $K^+$ .

Thin tissues can be imaged as a whole. In cases where a very detailed analysis on protein-mCLING colocalization is required, samples should be embedded in melamine and should be cut into thin slices (200 nm thick). If sample labeling is too dense, thickness can be reduced to 50 to 100 nm.

For efficient fixation of mCLING, a mixture of formaldehyde (FA) and glutaraldehyde is required. Scientists use self-made FA fixation solutions produced by dissolving paraformaldehyde (PFA) in phosphate buffered saline (PBS) or they apply a ready-to-use FA fixation solutions containing different amounts of methanol for stabilization.

#### Materials and reagents

- **Physiological buffer** (e.g. Hank's balanced salt solution or Jan & Jan's buffer depending on the tissue to be stained)
- **Physiological buffer** without  $Ca^{2+}$
- **mCLING (710 006AT488)** lyophilized
- **PBS:** Phosphate buffered saline, (200 mM NaCl, 2.5 mM KCl, 8 mM  $Na_2HPO_4$ , 1.5 mM  $KH_2PO_4$ ), pH 7.4
- **Fixation solution:** 4% FA, 0.2% glutaraldehyde in PBS
- **Ice**
- **Quenching buffer:** (100 mM  $NH_4Cl$ , 100 mM Glycine in PBS), pH 7.4
- **Primary antibody**
- **Secondary antibody:** Fluorophore conjugated species specific secondary antibody
- **Blocking buffer:** 0.5% Triton X-100, 10% normal serum in PBS (normal serum from the host-species of the secondary antibodies is recommended)
- **Incubation buffer:** 0.5% Triton X-100, 5% normal serum in PBS (normal serum from the host-species of the secondary antibodies is recommended)
- **Optional: DAPI nuclear stain**
- **Mounting medium**

#### Procedure

1. Dissect tissue using ice-cold **physiological buffer** without  $Ca^{2+}$ . Make sure to remove and clean muscles or cells of interest from other structures like connective, bone, and fat tissue. In poorly cleaned samples, mCLING binds to surrounding connective and fat tissue creating a strong background fluorescence signal.

2. Prepare two working **mCLING** solutions with a concentration between 2 to 1 nmol/ml, one with the standard **physiological buffer** and one with **physiological buffer without Ca<sup>2+</sup>**. For mammalian samples, keep solutions at 37°C for Drosophila larvae at room temperature (RT). Working mCLING concentrations must be determined via preliminary test experiments. Mammalian tissues require higher concentrations (sometimes up to 2 µM), as molecules are retained by the connective tissue layers lowering the effective concentration of the solution that reaches the cells of interest.

*Note: Avoid bright light when working with fluorescently labeled mCling to minimize photo bleaching of the fluorescent dye.*

3. Apply **mCLING** solution without Ca<sup>2+</sup> for 5 to 10 min (at 37°C for mammalian samples or at RT for Drosophila larvae experiments) to allow mCLING to penetrate into the tissue and to reach the membranes of the cell or structure of interest. Incubation times should also be established in preliminary test experiments. To determine optimal mCLING concentration and incubation time, perform parallel preparations and vary these two parameters. Samples can be embedded in melamine and cut into thin sections to be imaged in a conventional epifluorescence microscope. In this way, penetration of mCLING into the particular tissue can be evaluated.

4. Replace **mCLING** solution without Ca<sup>2+</sup> for the standard solution with mCLING and apply electrical stimulation of your choice. Alternatively, the standard solution can be replaced with a high K<sup>+</sup> solution containing the same concentration of mCLING. Briefly wash excess of mCLING with dye-free and Ca<sup>2+</sup>-free solution.

5. Immediately place the tissue in **fixation solution** and incubate for 30 min on ice. Incubate for another 30 min at RT.

6. Wash briefly with **PBS**.

7. Incubate for 30 min at RT in **quenching buffer**.

8. Wash briefly with **PBS**.

9. Incubate 3 times for 10 min each in **blocking buffer**.

10. Incubate with primary antibody diluted according to the manufacturer's instructions in **incubation buffer** for 1 h at RT. Overnight incubation is not recommended, as this might affect the stability of mCLING on the labeled membranes.

11. Wash 3 times for 10 min each with **incubation buffer**.

12. Incubate with **secondary antibody** diluted according to the manufacturer's instructions for 1 h at RT in **incubation buffer**.

*Optional: Add DAPI to the secondary antibody solution.*

**Notes:**

- *Avoid bright light when working with the secondary antibody to minimize photo bleaching of the fluorescent dye.*
- *In multiplex staining, make sure to use secondary antibodies cross-adsorbed against the host species of the other primary antibody used in your experiment. Ideally, all secondary antibodies should come from the same host species. If not, make sure that they have been cross-adsorbed against IgGs of the host-species of the other secondary antibody as well. This avoids cross-reaction between the secondary*

13. Wash 3 times for 10 min each with **PBS**.

14. Mount slides.

For more background information and more application specific protocols, refer to [Revelo NH and Rizzoli SO, 2016](#).